

Probiotic agents to protect the urogenital tract against infection¹⁻³

Gregor Reid

ABSTRACT The urogenital microflora of a healthy woman comprises ≈50 species of organisms, which differ in composition according to reproductive stages and exposure to several factors, including antibiotics and spermicides. Infections are very common with >300 million cases of urinary tract infections, bacterial vaginosis, and yeast vaginitis worldwide per annum. At the time of infection in the bladder and vagina, the urogenital flora is often dominated by the infecting pathogens, in contrast with healthy phases when indigenous organisms dominate. Premenopausal women have a flora of mostly lactobacilli, and certain properties of these strains, including adhesive ability and production of acids, bacteriocins, hydrogen peroxide, and biosurfactants, appear important in conferring protection to the host. Efforts to artificially restore an unbalanced flora with the use of probiotics have met with mixed results but research aimed at selecting scientifically based strains could well provide a reliable alternative treatment and preventive regimen to antibiotics in the future. *Am J Clin Nutr* 2001;73(suppl):437S–43S.

KEY WORDS Urogenital flora, infection, urinary tract, bacterial vaginosis, lactobacilli, probiotics

UROGENITAL INFECTIONS

Urogenital infections, defined here to include those that affect the bladder, kidneys, vagina, urethra, periurethra, and cervix, constitute a worldwide problem that affects >300 million women/y. In Western countries such as Canada and the United States, these infections are a common reason for a woman's visit to a family practitioner or urologist. For those infections in the urinary tract, it has been estimated that each episode leads to 6 d of signs and symptoms (1). The main clinical outcome is morbidity and discomfort amongst a large percentage of the female population, in addition to enormous costs to the health care system for treatment.

Many patients will experience a recurrence of symptoms, particularly within the first year of the original infection. Of more concern are infections that are complicated by kidney involvement because hospitalization is usually required and kidney damage or even death can occur. Serious complications can arise during pregnancy, sometimes leading to premature birth (2).

Historical data indicate that the vast majority (≈70%) of urinary tract infection (UTI) in a suburban, nonhospitalized community is caused by *Escherichia coli*, followed by other enterobacteriaceae and *Staphylococcus saprophyticus*. However, a recent 7 y study indicated a different result, whereby *E. coli*

infections were less common and *Enterococcus faecalis* was the second most prevalent uropathogen (3). The latter result was also found in hospitalized patients (4).

Most uncomplicated UTI cases are resolved between 1 and 7 d of antibiotic therapy. However, drug resistance to commonly used antibiotics (eg, trimethoprim/sulfamethoxazole) is increasing among uropathogens (3) and patients are experimenting more and more with alternative natural remedies such as cranberry juice, which appears to contain antiadhesive compounds that are active against uropathogens and can help prevent UTI (5–8).

Currently, preventive therapies for UTI rely almost exclusively on the use of antibiotics. In real terms, no true prophylaxis exists: current therapy involves long-term, low-dose antibiotic treatment, which involves the active killing of bacteria that enter the bladder.

Bacterial vaginosis (BV), formerly referred to as nonspecific vaginitis (9), is a major cause of urogenital disease in females. It has been defined as a mild infection of the lower female genital tract, characterized by the presence of 3 of 4 criteria to define urogenital disease: 1) release of an amine (putrescine, cadaverine, and trimethylamine) or fishy odor after the addition of 10% potassium hydroxide, 2) a vaginal pH >4.5, 3) clue cells in the vaginal fluid, and 4) a milky homogeneous vaginal discharge (10, 11). The clue-cell scoring of the cell population is termed as either normal (0–3; dominated by lactobacilli rods), intermediate (4–6; colonization by small gram-negative or gram-variable rods, eg, *Bacteroides* or *Gardnerella*), and curved gram-variable rods, eg, *Mobiluncus*), or BV (7–10; dominated by pathogens) (12).

The diagnosis of BV has its flaws (11, 13) and future improvements with the use of DNA probes, fast and accurate detection of amines, and short-chain volatile acids and enzymes should make diagnosis easier and more reliable for BV and other urogenital pathogens, eg, *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, herpes simplex virus, HIV, papillomavirus, *Treponema pallidum*, and *Trichomonas vaginalis* (14–18).

BV can lead to complications in pregnancies, causing premature rupture of the membranes, premature birth, or the death of the fetus or newborn (19–24). Premature rupture of the

¹From the Lawson Research Institute and the Departments of Microbiology and Immunology and Surgery (Urology), the University of Western Ontario, Canada.

²Presented at the symposium Probiotics and Prebiotics, held in Kiel, Germany, June 11–12, 1998.

³Address reprint requests to G Reid, H414, Lawson Health Research Institute, 268 Grosvenor Street, London, Ontario, N6A 4V2, Canada. E-mail: gregor@julian.uwo.ca.




FIGURE 1. When fimbriated, uropathogenic *Escherichia coli* enters the urogenital mucosa from the intestine, it may do so by spreading along the mucosa or by coming directly from the fluid phase into the bladder (A). It may also attach itself to the bare surface of a vaginal epithelial cell (B) or become part of the cell's bacterial biofilm (C), in which it either remains or becomes the dominant organism. When a patient has a bladder infection, the urogenital flora is invariably dominated by the infecting pathogen. The biofilm shown represents that from healthy premenopausal women. Lactobacilli (rods) dominate and coexist with gram-positive cocci (dark open and spiked circles representing capsule) and coliforms (clear, oval shape).

membranes can also be associated with BV, urinary tract infections, group B streptococcal infections, and the presence of organisms such as ureaplasma and mycoplasma in the urogenital tract (25–32).

DYNAMICS OF THE UROGENITAL MICROFLORA

The origin of the uropathogens in uncomplicated UTI and BV is the fecal flora. For >20 y, the key factor in pathogenesis has been regarded as the ability of the pathogens to attach to epithelial cells, thereby allowing them a niche in which to establish, multiply, spread, and avoid host defenses (33, 34). Numerous studies have investigated and documented the adhesins and receptor sites involved in the attachment process. Factors such as hemolysins, aerobactin, capsular antigens, and others play a role in *E. coli* pathogenesis in UTI (35–42). Furthermore, there are host susceptibility factors, including genetic determinants, age-related changes, and mucosal differences that influence the infection process (43).

Although the adhesion phenomenon is unquestionably important, there appears to be another key element of pathogenesis that is less well understood, namely the ability of the pathogens to survive exposure to the microflora that exists on the external urogenitalia. It has been reported that >50 species colonize the healthy vagina (44). Studies have shown that urogenital cells are covered by dense bacterial biofilms (45, 46) whose composition changes constantly but in which lactobacilli predominate, at least until menopause. Uropathogenic organisms emerge from the intestine (47) and come into contact with these biofilms on vaginal and urethral cells, yet little is known about what happens thereafter. It is presumed that uropathogens can either bypass the microflora or successfully enter the biofilms, survive, and continue their ascension into the bladder.

Biofilms have been described in the human host for >20 y (48, 49) but their role in health has not been studied as well as their association with disease. Current understanding indicates that the structure and dynamics of biofilms vary with the organisms, the surface to which they attach, the nutritional environment, and shear forces present in any given case. The organisms appear to benefit from biofilm formation by gaining access to nutrients,

escaping host immune cells and antimicrobial attack, and having an ability to better control their multiplication. A recurrent theme is that various factors, including nutrients, antimicrobials, and arriving bacteria, change the properties and the composition of the biofilms. In women, other factors such as hormonal concentrations, particularly estrogen, as well as changes induced by oral contraception, glycogen content, vaginal pH, steroid therapy, immunosuppression, and diseases (eg, diabetes mellitus) all influence the composition of the bacterial biofilms (50). The menstrual cycle appears to affect the adherence of lactobacilli to epithelial cells in healthy women in that days corresponding to high circulating concentrations of estrogens result in a higher adherence in vitro (51) and restored colonization postmenopause (52).

Studies have shown that antibiotic or spermicide exposure can cause disruption of the urogenital microflora and increase a woman's risk of infection (53, 54). Indeed, in vitro studies showed that most lactobacilli are eradicated by exposure to a low dose of nonoxynol-9, whereas uropathogens grow and prosper in high concentrations of this compound (55). Thus, protection of the host by her vaginal *Lactobacillus* flora is likely reduced upon exposure to spermicides. The fact that antimicrobials do not always completely wipe out the urogenital flora is due not only to resistance of individual strains, but also likely to the existence of biofilms, which themselves confer resistance to attack (56).

Scanning confocal laser microscopy has shown that biofilms are mushroom shaped and form water channels through which the organisms are nourished (57–59). With the use of such techniques, it should be possible to follow the entry of a uropathogen into the urogenital biofilm system, from whence it infects the host (Figure 1).

LACTOBACILLI AS GOOD PROBIOTICS FOR PREVENTION OF UROGENITAL INFECTIONS

The dominant presence of lactobacilli in the urogenital microflora of healthy women and the obliteration of lactobacilli in patients who develop UTI (60–62), BV, and many other genital infections (30, 63) [except candidiasis (64)] has led to a focus on these bacteria. Lactobacilli are gram-positive rods, primarily facultative or strict anaerobes that generally have a fastidious




FIGURE 2. Byproducts of lactobacillus metabolism that have an antagonistic effect against urinary and vaginal pathogens. The biosurfactants inhibit adhesion; the acids, bacteriocins, and hydrogen peroxide inhibit growth; and the coaggregation molecules block the spread of the pathogens.

growth requirement. They prefer an acidic environment and help create one by producing lactic and other acids. In general, lactobacilli have not been associated with disease and for >100 y have been regarded as nonpathogenic members of the intestinal and urogenital floras (65).

Lactobacilli have long been of interest to the dairy and agriculture industries (66, 67), although over the past century, studies in relation to human health were sporadic and often inconclusive. Some examples can be found in which lactic acid bacteria have been used to treat or prevent infections of the intestinal and genital tracts with different degrees of success (68–71). However, there has been failure in identifying the properties of lactobacilli required to prevent and treat disease; in determining the optimal dosage, duration, and mode of lactobacilli delivery; and including a placebo in the study. Studies have failed to recover and document lactobacilli properly and show their mechanisms of action.

A further problem with probiotic therapeutics is their reliability. In one study, this was highlighted by the fact that the contents of some health food products were inconsistent and different from those stated on the labels (72). More recently, we discovered that some health food products in Canada appear to contain ≥ 1 common *Lactobacillus* strain, apparently identical to ATCC strain 21052, and that US and Canadian products contain *Lactobacillus plantarum*, although its presence is not stated on the food label (73).

In more recent years, the use of probiotics per se and lactobacilli specifically has received greater attention as an alternative, inexpensive, and natural remedy to restore and maintain health (74–79). Two strains, *Lactobacillus GG* (ATCC 53103) and *Lactobacillus rhamnosus GR-1* appear to be effective at colonizing and protecting the intestine (80–86) and urogenital tract (87–91), respectively, against microbial infection.

There is one obvious question: what properties do these strains possess that make them effective probiotic agents? The answer is not fully known, but some common denominators appear to exist, namely the ability to adhere to and colonize tissues and the capacity to inhibit the pathogenesis of disease-causing organisms (Figure 2). Another question can be raised: do we expect an exogenous probiotic strain to colonize the gastrointestinal and urogenital tracts of a given person for a long time and even become part of the normal flora, replacing or coexisting with the endogenous lactobacilli organisms? Or is the aim to

substitute the endogenous bacteria only while the normal flora is repressed (eg, as a result of antibiotic therapy) (92)? Again, no clear answers are available.

To more fully ascertain which properties are required by lactobacilli to protect the host, we recommend a series of extensive microbiological, physico-chemical, and molecular biology methodologies (89, 93–97). It appears that a given strain of *Lactobacillus* can express several, but not necessarily all, of the known key factors and be able to compete in the urogenital microenvironment. For example, lactobacilli can use many mechanisms to adhere to surfaces, such as electrostatic, hydrophobic, hydrophilic, capsular, and fimbrial mechanisms (95, 98–102): in the urogenital tract, hydrophilic *L. rhamnosus GR-1* and hydrophobic *Lactobacillus fermentum B-54* both colonize (91). Some strains can bind better to intestinal cells and inhibit pathogen adhesion (103, 104), but they may not be able to effectively inhibit growth of uropathogens (94).

Before commencing more human studies, *L. rhamnosus GR-1* was found to be the best of a group of 34 *Lactobacillus* strains isolated from dairy, poultry, health food, and human sources, with respect to adhesion to squamous and transitional uroepithelial cells, competitive exclusion of pathogens, and production of inhibitors of uropathogen growth (94). In 1995 another characteristic of lactobacilli was discovered that appeared to be important in conferring probiotic action against uropathogens. Fifteen strains were found to produce biosurfactant (105). The substance or substances adsorbed to surfaces and inhibited the initial adhesion of *E. faecalis* by 70%. The crude substance was analyzed and was found to contain proteins and carbohydrates. Amino acid analysis of hydrolyzed material from strain *Lactobacillus acidophilus RC-14* showed a high alanine content compared with less active strains. The activity is not due to lipoteichoic acid or glycosyldiglycerides or to factors such as acid or bacteriocins, which inhibit bacterial growth. Sodium dodecyl sulphate-polyacrylamide gel electrophoresis showed a variety of proteins with molecular masses of from 14.4 to 140 kDa (106, 107). The biosurfactant activity is resistant to trypsin and pepsin, and sensitive to α -amylase and lysozyme, and resistant to 75°C degree heating. The antiadhesive molecules produced by certain lactobacilli hold promise for application to many human sites where pathogens attach, colonize, and confer disease. More recently, the activity was shown to affect a broad range of pathogens (Table 1) (108) and an active component was found to be a collagen binding protein (109). The

TABLE 1
Pathogens affected by biosurfactant activity¹

Strain	Adhesion (4 h) $\times 10^6 \text{ cm}^{-2}$
<i>Escherichia coli</i> 67	
No biosurfactant	2.4
With biosurfactant	0.5
<i>E. coli</i> Hu734	
No biosurfactant	1.1
With biosurfactant	0.2
<i>Enterococcus faecalis</i> 1131	
No biosurfactant	1.04
With biosurfactant	0.04
<i>E. faecalis</i> 1396	
No biosurfactant	0.3
With biosurfactant	0.02
<i>Klebsiella pneumoniae</i> 3a	
No biosurfactant	0.9
With biosurfactant	0.7
<i>Proteus mirabilis</i> 28cii	
No biosurfactant	1.0
With biosurfactant	0.5
<i>Providencia stuartii</i> UHL103	
No biosurfactant	8.9
With biosurfactant	4.0
<i>Pseudomonas aeruginosa</i> AK1	
No biosurfactant	11.0
With biosurfactant	9.5
<i>Staphylococcus epidermidis</i> 3059	
No biosurfactant	1.8
With biosurfactant	0.2

¹See reference 104.

discovery of biosurfactants in lactobacilli and several antiadhesion components is an exciting development.

Hydrogen peroxide-producing strains are believed to be important in vaginal colonization (110), yet *L. rhamnosus* GR-1, which colonizes the vagina is a nonproducer. Perhaps its resistance to spermicide is a more important attribute for its probiotic usage because these agents can destroy the normal flora (55, 111, 112). Lactic acid was shown to inhibit growth of *Gardnerella vaginalis* more so than hydrogen peroxide, unless the latter is in the presence of myeloperoxidase (113). Strains with identical molecular profiles, eg, poultry isolate *L. rhamnosus* A-60 and vaginal isolate *L. rhamnosus* RC-17 (73), may actually utilize quite different mechanisms for colonization (namely high adhesiveness or production of inhibitory substances for *L. rhamnosus* RC-17 and high capacity to exclude pathogens in the case of *L. rhamnosus* A-60). *L. rhamnosus* GR-1 and strain 36 have similar molecular profiles, yet the former produces a biosurfactant active in preventing enterococci from adhering to surfaces (105), whereas the latter has a biosurfactant that is inactive.

Presumably, lactobacilli continually enter the intestine from food sources. There are studies that indicate that certain species, eg, *L. acidophilus*, are better able to survive stomach pH and bile salt exposure so that they may pass through into the intestine (114). In other studies of biopsied intestinal mucosa, strains of *L. plantarum*, *Lactobacillus casei* subsp. *rhamnosus*, *Lactobacillus reuteri*, and *Lactobacillus agilis* were recovered postimplantation (115). However, without the use of molecular typing and specific probes, one cannot be certain that these strains are transmitted to, or subsequently colonize, the female urogenital tract. One could

argue that the strains that appear in stool and in vaginal mucosa throughout life are those that colonized the urogenital tract shortly after birth, and that subsequent events, eg, nutritional and hormonal changes, led to the emergence or suppression of certain strains. A lengthy and large epidemiologic study might be needed to better understand the succession of the normal flora during different phases of life.

There is an indication that probiotic strains can undergo genetic changes during colonization: a tetracycline-sensitive *Bifidobacterium* strain administered to antibiotic-decontaminated mice maintained in isolators was found to produce 2 undistinguished intestinal populations according to restriction endonuclease patterns of total DNA, but one variant was highly resistant to tetracycline (V Smeianov et al, unpublished observations, 1998). Both variants coexisted in the mouse intestine and there was a strong advantage of the newly appeared variant. This possibility of genetic adaptation or in vivo selection of probiotic bacteria is an area of interest for our group.


We assume that artificial colonization is possible and that in vitro analyses will allow us to construct methods for selecting and delivering these strains. However, an adjunct to such studies must be an examination of the strains already colonizing healthy humans and a clarification of how these organisms are altered by food intake. On the basis of our knowledge, strains with different origins and probiotic properties seem to exist in the vagina of healthy women (73, 116). The challenge will be to identify which strains are the most beneficial for health and why.

Further proof of urogenital colonization and protection from infection was obtained from a clinical trial in which 55 premenopausal women were given weekly either one suppository of *L. rhamnosus* GR-1 and B-54 (0.5 g) or one suppository of a lactobacillus growth factor for 1 y. The patients were followed up after 2 wk and then monthly (91). Six patients were excluded in the first month for noncompliance, moving, or becoming pregnant, and 11 patients did not complete the study in full; however, their data were included to examine infection rates. Again, there were no side effects in the study. The UTI infection rate decreased from 6.0 per previous year to 1.6 (73% decrease) for those given lactobacilli and to 1.3 (79% decrease) for those given *Lactobacillus* growth factor. Time to first infection was the same in both groups, namely 21–22 wk. Vaginal pH was the same for both groups, namely 4.6–5.0. The viable lactobacillus counts recovered from vaginal swabs increased with therapy, especially for months 7–12 for lactobacilli-treated patients, during which time lower UTI rates were seen.

It had been hypothesized by some researchers that a low vaginal pH (<5) was sufficient to prevent infection, including UTI, sexually transmitted diseases, and AIDS. An analysis of 10 patients showed that lactobacillus therapy maintained the vaginal pH at acidic levels (4.8 ± 0.5) during the study (117). The UTI rates fell by 79.4% (94) but UTI did occur despite a vaginal pH of 4.8 and the causative agents were common uropathogens (6 *E. coli*, 3 streptococci, and 3 staphylococci). Thus, from these studies (91, 117), it was concluded that a vaginal acid pH < 5 was not sufficient alone to prevent UTI.

CONCLUSIONS

There is now growing evidence that certain species and strains present in the healthy urogenital tract protect the host against

infection by pathogenic microorganisms. Many properties required to confer this protection have been identified, but evidence of their expression in vivo is scant and the relative significance of each is unknown. The concept of treating and preventing urogenital infection by instilling probiotic organisms has great appeal to patients and caregivers. The ability to administer *L. rhamnosus* GR-1 and *L. fermentum* RC-14, which colonize the intestine and vagina, by mouth to help restore and maintain urogenital health (118) provides a major step in the right direction for patients as it potentially allows for the self administration of therapy. However, for this approach to be successful, proper selection of strains, proof of concept, and efficacy must accompany products used in patients. 

REFERENCES

1. Foxman B, Fredrichs RR. Epidemiology of UTI. Diaphragm use and sexual intercourse. *Am J Public Health* 1985;75:1308-13.
2. Bergeron MG. Treatment of pyelonephritis in adults. *Med Clin N Am* 1972;79:619-49.
3. Reid G, Seidenfeld A. Drug resistance amongst uropathogens isolated from women in a suburban population: 7 years of change. *Can J Urol* 1997;4:432-7.
4. Preston CAK, Bruce AW, Reid G. Antibiotic resistance of urinary pathogens isolated from patients attending The Toronto Hospital between 1986 and 1990. *J Hosp Infect* 1992;21:129-35.
5. Sobota AE. Inhibition of bacterial adherence by cranberry juice: potential use for the treatment of urinary tract infection. *J Urol* 1984;131:1013-6.
6. Schmidt DR, Sobota AE. An examination of the anti-adherence activity of cranberry juice on urinary and non-urinary bacterial isolates. *Microbios* 1988;55:171-81.
7. Avorn J, Mohane M, Gurwitz JH, Glynn RJ, Choordnovsky I, Lipsitz LA. Reduction of bacteriuria after ingestion of cranberry juice. *JAMA* 1994;271:751-4.
8. Ofek I, Goldhar J, Sharon N. Anti-*Escherichia coli* adhesin activity of cranberry and blueberry juices. *Adv Exp Med Biol* 1996;408:179-83.
9. Gardner HL, Dukes CD. *Haemophilus vaginalis* vaginitis: a newly defined specific infection previously classified "non-specific" vaginitis. *Am J Obstet Gynecol* 1955;69:962-76.
10. Amsel R, Totten PA, Spiegel CA, Chen KCS, Eschenbach D, Holmes KK. Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med* 1983;74:14-22.
11. Hillier SL. Diagnostic microbiology of bacterial vaginosis. *Am J Obstet Gynecol* 1993;169:455-9.
12. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of Gram stain interpretation. *J Clin Microbiol* 1991;29:297-301.
13. Hay PE, Taylor-Robinson D. Defining bacterial vaginosis: to BV or not to BV, that is the question. *Int J STD AIDS* 1996;7:233-5.
14. Aynaud O, Bijaoui G, Huynh B. Genital bacterial infections associated with papillomavirus: value of screening and basis for treatment. *Contracept Fertil Sex* 1993;21:149-52.
15. Graves A, Gardner WA Jr. Pathogenicity of *Trichomonas vaginalis*. *Clin Obstet Gynecol* 1993;36:145-52.
16. Biro FM, Rosenthal SL, Kinyalocots M. Gonococcal and chlamydia genitourinary infections in symptomatic and asymptomatic adolescent women. *Clin Pediatr* 1995;34:419-23.
17. Nilsson U, Hellerg D, Shoubnikova M, Nilsson S, Mardh PA. Sexual behavior risk factors associated with bacterial vaginosis and *Chlamydia trachomatis* infection. *Sex Transm Dis* 1997;24:241-6.
18. Sewankambo N, Gray RH, Wawer MJ, et al. HIV-1 infection associated with abnormal vaginal flora morphology and bacterial vaginosis. *Lancet* 1997;350:530-1.
19. Alger L, Pupkin M. Etiology of preterm rupture of membranes. *Clin Obstet Gynecol* 1986;29:758-70.
20. Gibbs RS, Romero R, Hillier SL, Eschenbach DA, Sweet RL. A review of premature birth and subclinical infection. *Am J Obstet Gynecol* 1992;166:1515-28.
21. McGregor JA, French JI, Seo K. Premature rupture of membranes and bacterial vaginosis. *Am J Obstet Gynecol* 1993;169:463-6.
22. Chaim W, Mazor M, Leiberman JR. The relationship between bacterial vaginosis and preterm birth. *Arch Gynecol Obstet* 1997;259:51-8.
23. Goldenberg RL, Andrews WW, Yuan AC, MacKay HT, St. Louis ME. Sexually transmitted diseases and adverse outcomes of pregnancy. *Clin Perinatol* 1997;24:23-41.
24. Newton ER, Piper J, Peairs W. Bacterial vaginosis and intraamniotic infection. *Am J Obstet Gynecol* 1997;176:672-7.
25. Kass EH. Bacteriuria and pyelonephritis of pregnancy. *Arch Intern Med* 1960;105:194-8.
26. Braun P, Lee YH, Klein JO, et al. Birth weight and genital mycoplasmas in pregnancy. *N Engl J Med* 1971;284:167-71.
27. Brumfitt W. The effects of bacteriuria in pregnancy on maternal and fetal health. *Kidney Int Suppl* 1975;4:113-9.
28. Regan JA, Chao S, James LS. Premature rupture of membranes, preterm delivery and group B streptococcal colonization of mothers. *Am J Obstet Gynecol* 1981;141:184-6.
29. Eschenbach DA, Nugent RP, Rao AV, et al. A randomized placebo-controlled trial of erythromycin for the treatment of *Ureaplasma urealyticum* to prevent premature delivery. *Am J Obstet Gynecol* 1991;164:734-42.
30. Hay PE, Lamont RF, Taylor-Robinson D, Morgan J, Ison C, Pearson J. Abnormal bacterial colonisation of the genital tract and subsequent preterm delivery and late miscarriage. *BMJ* 1994;308:295-8.
31. Holst E, Goffeng AR, Andersch B. Bacterial vaginosis and vaginal microorganisms in idiopathic premature labor and association with pregnancy outcome. *J Clin Microbiol* 1994;32:176-86.
32. Calleri L, Porcelli A, Gallelo D, Taccani C, Surico N. Bacterial vaginosis and premature membrane rupture: an open study. *Minerva Ginecol* 1997;49:19-23.
33. Eden CS, Hanson LA, Jodal U, Lindberg U, Akerlund AS. Variable adherence to normal human urinary-tract epithelial cells of *Escherichia coli* strains associated with various forms of urinary-tract infection. *Lancet* 1976;1:490-2.
34. Reid G, Sobel JD. Bacterial adherence in the pathogenesis of urinary tract infection: a review. *Rev Infect Dis* 1987;9:470-87.
35. Brooks HJL, O'Grady F, McSherry MA, Cattell MR. Uropathogenic properties of *Escherichia coli* in recurrent urinary-tract infection. *J Med Microbiol* 1980;13:57-68.
36. Hacker J, Schmidt G, Hughes C, Knapp S, Marget M, Goebel W. Cloning and characterization of genes involved in production of mannose-resistant, neuraminidase-susceptible (X) fimbriae from a uropathogenic 06:K15:H31 *Escherichia coli* strain. *Infect Immun* 1985;47:434-40.
37. Hultgren SJ, Porter TN, Schaeffer AJ, Duncan JL. Role of type 1 pili and effects of phase variation on lower urinary tract infections produced by *Escherichia coli*. *Infect Immun* 1985;50:370-7.
38. Kallenius G, Molby R, Winberg J. Adhesion of *Escherichia coli* to periurethral cells correlated to mannose-resistant agglutination of human erythrocytes. *FEMS Microbiol Lett* 1979;5:295-9.
39. Kist ML, Salit IE, Hofmann T. Purification and characterization of the Dr hemagglutinins expressed by two uropathogenic *Escherichia coli* strains. *Infect Immun* 1990;58:695-702.
40. Pere A, Leinonen M, Vaisanen-Rhen V, Rhen M, Korhonen TK. Occurrence of type-1C fimbriae on *Escherichia coli* strains isolated from human extraintestinal infections. *J Gen Microbiol* 1985;131:1705-11.
41. Rhen M, Klemm P, Korhonen TK. Identification of two new hemagglutinins of *Escherichia coli*, N-acetyl-D-glucosamine-specific fimbriae and a blood group M-specific agglutinin, by cloning the corresponding genes in *Escherichia coli* K-12. *J Bacteriol* 1986;168:1234-42.



42. Vaisanen V, Korhonen TK, Jokinen M, Gahmberg CG, Ehnholm C. Blood group M specific haemagglutinin in pyelonephritogenic *Escherichia coli*. Lancet 1982;1:1192 (letter).
43. Scholes D, Hooton TM, Roberts PL, Stapleton AE, Gupta K, Stamm WE. Risk factors for recurrent urinary tract infection in young women. J Infect Dis 2000;182:1177–82.
44. Redondo-Lopez V, Cook RL, Sobel JD. Emerging role of lactobacilli in the control and maintenance of the vaginal bacterial microflora. Rev Infect Dis 1990;12:856–72.
45. Sadhu K, Domingue PAG, Chow AW, Nelligan J, Bartlett K, Costerton JW. A morphological study of the *in situ* tissue-associated autochthonous microflora of the human vagina. Microb Ecol Health Dis 1989;2:99–106.
46. Reid G, McGroarty JA, Domingue PAG, et al. Coaggregation of urogenital bacteria *in vitro* and *in vivo*. Curr Microbiol 1990; 20: 47–52.
47. Wold AE, Caugant DA, Lidin-Janson G, de Man P, Svanborg C. Resident colonic *Escherichia coli* strains frequently display uropathogenic characteristics. J Infect Dis 1992;165:46–52.
48. Savage DC. Microbial ecology of the gastrointestinal tract. Annu Rev Microbiol 1977;31:107–33.
49. Costerton JW, Irwin RT, Cheng K-J. The bacterial glycocalyx in nature and disease. Annu Rev Microbiol 1981;35:399–424.
50. Galask RP. Vaginal colonization by bacteria and yeast. Am J Obstet Gynecol 1988;158:993–5.
51. Chan RCY, Bruce AW, Reid G. Adherence of cervical, vaginal and distal urethral normal microbial flora to human uroepithelial cells and the inhibition of adherence of uropathogens by competitive exclusion. J Urol 1984;131:596–601.
52. Raz R, Stamm WE. A controlled trial of intravaginal estriol in postmenopausal women with recurrent urinary tract infections. N Engl J Med 1993;329:753–6.
53. Hooton TM, Hillier S, Johnson C, Roberts PL, Stamm WE. *Escherichia coli* bacteriuria and contraceptive method. JAMA 1991; 265:64–9.
54. Reid G, Bruce AW, Cook RL, Llano M. Effect on the urogenital flora of antibiotic therapy for urinary tract infection. Scand J Infect Dis 1990;22:43–7.
55. McGroarty JA, Chong S, Reid G, Bruce AW. Influence of the spermicidal compound nonoxonyl-9 on the growth and adhesion of urogenital bacteria *in vitro*. Curr Microbiol 1990;21:219–23.
56. Costerton JW, Lappin-Scott HM. Introduction to microbial biofilms. In: Lappin-Scott HM, Costerton JW, eds. Microbial biofilms. Cambridge, UK: Cambridge University Press, 1995:1–11.
57. James GA, Korber DR, Caldwell DE, Costerton JW. Digital image analysis of growth and starvation responses of surface-colonizing *Actinobacter* sp. J Bacteriol 1995;177:907–15.
58. Sanford BA, de Feijter AW, Wade MH, Thomas VL. A dual fluorescence technique for visualization of *Staphylococcus epidermidis* biofilm using scanning confocal laser microscopy. J Industr Microbiol 1996;16:48–56.
59. Potera C. Biofilms invade microbiology. Science 1996;273:1795–7.
60. Stamey TA. The role of introital enterobacteria in recurrent urinary infections. J Urol 1973;109:467–72.
61. Schaeffer AJ, Stamey TA. Studies of introital colonization in women with recurrent urinary infections. IX. The role of antimicrobial therapy. J Urol 1977;118:221–4.
62. Seddon JM, Bruce AW, Chadwick P, Carter D. Introital bacterial flora—effect of increased frequency of micturition. Br J Urol 1976; 48:211–8.
63. Hillier SL, Krohn MA, Rabe LK, Klebanoff SJ, Eschenbach DA. The normal vaginal flora, H₂O₂-producing lactobacilli, and bacterial vaginosis in pregnant women. Clin Infect Dis 1993;16(suppl):S273–81.
64. Sobel JD. Biotherapeutic agents as therapy for vaginitis. In: Elmer GW, MacFarland K, Surawicz C, eds. Biotherapeutic agents and infectious diseases. Totowa, NJ: Humana Press: 221–44.
65. Bibel DJ. Elie Metchnikoff's bacillus of long life. ASM News 1988;54:661–5.
66. Klaenhammer TR. Microbiological considerations in selection and preparation of *Lactobacillus* strains for use as dietary adjuncts. J Dairy Sci 1982;65:1339–49.
67. Watkins BA, Miller BF, Neil DH. *In vivo* inhibitory effects of *Lactobacillus acidophilus* against pathogenic *Escherichia coli* in gnotobiotic chicks. Poult Sci 1982;61:1298–308.
68. Beck C, Necheles H. Beneficial effects of administration of *Lactobacillus acidophilus* in diarrheal and other intestinal disorders. Am J Gastroenterol 1961;35:522–30.
69. Alexander JG. Thrush bowel infection: existence, incidence, prevention and treatment, particularly by a *Lactobacillus acidophilus* preparation. Curr Med Drugs 1967;8:3–11.
70. Clements ML, Levine MM, Black RE, et al. *Lactobacillus* prophylaxis for diarrhea due to enterotoxigenic *Escherichia coli*. Antimicrob Agents Chemother 1981;20:104–8.
71. Hilton E, Isenberg HD, Alperstein P, France K, Borenstein MT. Ingestion of yogurt containing *Lactobacillus acidophilus* as prophylaxis for candidal vaginitis. Ann Intern Med 1992;116:353–7.
72. Hughes VL, Hillier SL. Microbiologic characteristics of *Lactobacillus* products used for colonization of the vagina. Obstet Gynecol 1990;75:244–8.
73. Zhong W, Millsap K, Bialkowska-Hobrzanska H, Reid G. Differentiation of *Lactobacillus* species by molecular typing. Appl Environ Microbiol 1998;64:2418–23.
74. McGroarty JA. Probiotic use of lactobacilli in the human female urogenital tract. FEMS Immunol Med Microbiol 1993;6:251–64.
75. Chang H. Genetic engineering to enhance microbial interference and related therapeutic applications. Nat Biotechnol 1996;14:444–7.
76. Elmer GW, Surawicz CM, McFarland LV. Biotherapeutic agents: a neglected modality for the treatment and prevention of selected intestinal and vaginal infections. JAMA 1996;275:870–6.
77. Reid G. Probiotics: how microorganisms compete. JAMA 1996;276: 29–30.
78. Shalev E, Battino S, Weiner E, Colodner R, Keness Y. Ingestion of yogurt containing *Lactobacillus acidophilus* compared with pasteurized yogurt as prophylaxis for recurrent candidal vaginitis and bacterial vaginosis. Arch Fam Med 1996;5:593–6.
79. Hamilton-Miller JMT. Living in the 'post-antibiotic era': could the use of probiotics be an effective strategy? Clin Microbiol Infect 1997;3:2–3.
80. Biller JA, Katz AJ, Flores AF, Buie TM, Gorbach SL. Treatment of recurrent *Clostridium difficile* colitis with *Lactobacillus* GG. J Pediatr Gastroenterol Nutr 1995;21:224–6.
81. Gorbach SL, Chang TW, Goldin B. Successful treatment of relapsing *Clostridium difficile* colitis with *Lactobacillus* GG. Lancet 1987;2:1519 (letter).
82. Isolauri E, Juntunen M, Rautanen T. A human *Lactobacillus* strain (*Lactobacillus* GG) promotes recovery from acute diarrhea in children. Pediatrics 1991;88:90–7.
83. Oksanen P, Salminen S, Saxelin M, et al. Prevention of traveler's diarrhea by *Lactobacillus* GG. Ann Med 1990;22:53–6.
84. Siitonen S, Vapaatalo H, Salminen S, et al. Effect of *Lactobacillus* GG yogurt and prevention of antibiotic-associated diarrhea. Ann Med 1990;22:57–9.
85. Saxelin M, Pessi T, Salminen S. Fecal recovery following oral administration of *Lactobacillus* strain GG (ATCC 53103) in gelatine capsules to healthy volunteers. Int J Food Microbiol 1995;25:199–203.
86. Miller MR, Bacon C, Smith SL, Walker V, Hall MA. Enteral feeding of premature infants with *Lactobacillus* GG. Arch Dis Child 1993;69:483–7.
87. Bruce AW, Reid G. Intravaginal instillation of lactobacilli for prevention of recurrent urinary tract infections. Can J Microbiol 1988;34:339–43.
88. Bruce AW, Reid G, McGroarty JA, Taylor M, Preston C. Preliminary study on the prevention of recurrent urinary tract infections in ten adult women using intravaginal lactobacilli. Int Urogynecol J 1992;3:22–5.



89. Reid G, Millsap K, Bruce AW. Implantation of *Lactobacillus casei* var *rhamnosus* into vagina. *Lancet* 1994;344:1229 (letter).
90. Hilton E, Rindos P, Isenberg HD. *Lactobacillus* GG vaginal suppositories and vaginitis. *J Clin Microbiol* 1995;33:1433 (letter).
91. Reid G, Bruce AW, Taylor M. Instillation of *Lactobacillus* and stimulation of indigenous organisms to prevent recurrence of urinary tract infections. *Microecol Ther* 1995;23:32-45.
92. Korshunov VM, Sinitsyna NA, Ginodman GA, Pinegin BV. Correction of the intestinal microflora in cases of chemotherapeutic dysbacteriosis by means of bifidobacterial and lactobacterial autostrains. *Zh Mikrobiol Epidemiol Immunobiol* 1985;9:20-5.
93. Geertsema-Doornbusch GI, Noordmans J, Bruce AW, et al. Microelectrophoresis as a method to study heterogeneity of microbial cell surfaces: agreement between microelectrophoresis and electron microscopy applied to lactobacilli after serial passaging. *J Microbiol Methods* 1994;19:269-77.
94. Reid G, Cook RL, Bruce AW. Examination of strains of lactobacilli for properties which may influence bacterial interference in the urinary tract. *J Urol* 1987;138:330-5.
95. Reid G, Cuperus PL, Bruce AW, et al. Comparison of contact angles and adhesion to hexadecane of urogenital, dairy and poultry lactobacilli: effect of serial culture passages. *Appl Environ Microbiol* 1992;58:1549-53.
96. Millsap K, Reid G, van der Mei HC, Busscher HJ. Adhesion and displacement of *Enterococcus faecalis* by *Lactobacillus* and *Streptococcus* sp. from hydrophobic and hydrophilic substrate as studied in a parallel plate flow chamber. *Appl Environ Microbiol* 1994;60:1867-74.
97. Millsap K, Reid G, van der Mei HC, Busscher HJ. Cluster analysis of genotypically characterized *Lactobacillus* species based on physicochemical cell surface properties and their adhesion to hexadecane. *Can J Microbiol* 1997;43:284-91.
98. Cook RL, Harris RJ, Reid G. Effect of culture media and growth phase on the morphology of lactobacilli and on their ability to adhere to epithelial cells. *Curr Microbiol* 1988;17:159-66.
99. Cuperus P, van der Mei HC, Reid G, et al. The effect of serial passaging of lactobacilli in liquid medium on their physico-chemical and structural surface characteristics. *Cell Material* 1992;2:271-80.
100. Cuperus PL, van der Mei HC, Reid G, et al. Physico-chemical surface characteristics of urogenital and poultry lactobacilli. *J Colloid Interface Sci* 1993;156:319-24.
101. McGroarty JA. Cell surface appendages of lactobacilli. *FEMS Microbiol Lett* 1994;124:405-10.
102. Andreu A, Stapleton AE, Fennell CL, Hillier SL, Stamm WE. Hemagglutination, adherence, and surface properties of vaginal *Lactobacillus* species. *J Infect Dis* 1995;171:1237-43.
103. Reid G, Servin A, Bruce AW, Busscher HJ. Adhesion of three *Lactobacillus* strains to human urinary and intestinal epithelial cells. *Microbios* 1993;75:57-65.
104. Hudault S, Bridonneau C, Raibaud P, Chabanet C, Vial MF. Relationship between intestinal colonization of *Bifidobacterium bifidum* in infants and the presence of exogenous and endogenous growth-promoting factors in their stools. *Pediatr Res* 1994;35:696-700.
105. Velraeds MC, van der Mei HC, Reid G, Busscher HJ. Inhibition of initial adhesion of uropathogenic *Enterococcus faecalis* by biosurfactants from *Lactobacillus* isolates. *Appl Environ Microbiol* 1996;62:1958-63.
106. Velraeds MM, van der Mei HC, Reid G, Busscher HJ. Physico-chemical and biochemical characterization of biosurfactants released from *Lactobacillus* strains. *Colloids Surfaces B: Biointerfaces* 1996;8:51-61.
107. Velraeds MM, van der Mei HC, Reid G, Busscher HJ. Inhibition of initial adhesion of uropathogenic *Enterococcus faecalis* to solid substrate by an adsorbed biosurfactant layer from *Lactobacillus acidophilus*. *Urology* 1997;49:790-4.
108. Velraeds MC, van der Belt B, van der Mei HC, Reid G, Busscher HJ. Interference in initial adhesion of uropathogenic bacteria and yeasts to silicone rubber by a *Lactobacillus* biosurfactant. *J Med Microbiol* 1998;47:1081-5.
109. Heinemann C, van Hylckama Vlieg JE, Janssen DB, Busscher HJ, van der Mei HC, Reid G. Purification and characterization of a surface-binding protein from *Lactobacillus fermentum* RC-14 that inhibits adhesion of *Enterococcus faecalis* 1131. *FEMS Microbiol Lett* 2000;190:177-180.
110. Eschenbach DA, Davick PR, Williams SJ, Klebanoff SJ, Young-Smith CM, Holmes KK. Prevalence of hydrogen peroxide-producing *Lactobacillus* species in normal women and women with bacterial vaginosis. *J Clin Microbiol* 1989;27:251-6.
111. McGroarty JA, Soboh F, Bruce AW, Reid G. The spermicidal compound nonoxynol-9 increases adhesion of *Candida* species to human epithelial cells in vitro. *Infect Immun* 1990;58:2005-7.
112. McGroarty JA, Tomeczek L, Pond DG, Reid G, Bruce AW. Hydrogen peroxide production by *Lactobacillus* species, correlation with susceptibility to the spermicidal compound nonoxynol-9. *J Infect Dis* 1992;165:1142-4.
113. McLean NW, McGroarty JA. Growth inhibition of metronidazole-susceptible and metronidazole-resistant strains of *Gardnerella vaginalis* by lactobacilli in vitro. *Appl Environ Microbiol* 1996;62:1089-92.
114. Sarra PC, Dellaglio F. Colonization of a human intestine by four different genotypes of *Lactobacillus acidophilus*. *Microbiologia* 1984;7:331-9.
115. Johansson M-L, Molin G, Jeppsson B, Nobaek S, Ahrne S, Bengmark S. Administration of different *Lactobacillus* strains in fermented oatmeal soup: in vivo colonization of human intestinal mucosa and effect on the indigenous flora. *Appl Environ Microbiol* 1993;59:15-20.
116. Reid G, McGroarty JA, Tomeczek L, Bruce AW. Identification and plasmid profiles of *Lactobacillus* species from the vagina of 100 women. *FEMS Immunol Med Microbiol* 1996;15:23-6.
117. Reid G, Bruce AW. Low vaginal pH and urinary-tract infection. *Lancet* 1995;346:1704 (letter).
118. Reid G, Bruce AW, Fraser N, Heinemann C, Owen J, Henning B. Oral probiotics can resolve urogenital infections. *FEMS Med Microbiol Immunol* 2000 (in press).

