

Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies¹⁻⁴

Gary Williamson and Claudine Manach

ABSTRACT

For some classes of dietary polyphenols, there are now sufficient intervention studies to indicate the type and magnitude of effects among humans *in vivo*, on the basis of short-term changes in biomarkers. Isoflavones (genistein and daidzein, found in soy) have significant effects on bone health among postmenopausal women, together with some weak hormonal effects. Monomeric catechins (found at especially high concentrations in tea) have effects on plasma antioxidant biomarkers and energy metabolism. Procyanidins (oligomeric catechins found at high concentrations in red wine, grapes, cocoa, cranberries, apples, and some supplements such as Pycnogenol) have pronounced effects on the vascular system, including but not limited to plasma antioxidant activity. Quercetin (the main representative of the flavonol class, found at high concentrations in onions, apples, red wine, broccoli, tea, and *Ginkgo biloba*) influences some carcinogenesis markers and has small effects on plasma antioxidant biomarkers *in vivo*, although some studies failed to find this effect. Compared with the effects of polyphenols *in vitro*, the effects *in vivo*, although significant, are more limited. The reasons for this are 1) lack of validated *in vivo* biomarkers, especially in the area of carcinogenesis; 2) lack of long-term studies; and 3) lack of understanding or consideration of bioavailability in the *in vitro* studies, which are subsequently used for the design of *in vivo* experiments. It is time to rethink the design of *in vitro* and *in vivo* studies, so that these issues are carefully considered. The length of human intervention studies should be increased, to more closely reflect the long-term dietary consumption of polyphenols. *Am J Clin Nutr* 2005;81(suppl):243S–55S.

KEY WORDS Polyphenols, flavonoids, procyanidin, bioavailability, isoflavone, quercetin, catechin

INTRODUCTION

It is clear that food components must, by definition, be bioavailable in some form to exert biological effects. There have been major advances in the past few years in our knowledge regarding polyphenol absorption and metabolism (1–4), and it is apparent that most classes of polyphenols are sufficiently absorbed to have the potential to exert biological effects. For example, quercetin after a meal containing onions, catechins after red wine consumption, and isoflavones after soy consumption reach micromolar concentrations in the blood (1, 2, 5–7). These findings demonstrate that polyphenols cross the intestinal barrier and reach concentrations in the bloodstream that have been shown to exert effects *in vitro*, in some studies.

There are thousands of articles on the effects of polyphenols on biological systems *in vitro*. However, many of those studies did

not take bioavailability and metabolism factors into consideration, and the effects reported in those studies do not necessarily occur *in vivo*. Although most polyphenols are absorbed to some extent, this is very dependent on the type of polyphenol. The range of concentrations required for an effect *in vitro* varies from <0.1 $\mu\text{mol/L}$ to >100 $\mu\text{mol/L}$. Because physiologic concentrations do not exceed 10 $\mu\text{mol/L}$, the effects of polyphenols *in vitro* at concentrations of >10 $\mu\text{mol/L}$ are generally not valid, with the possible (but unproven) exception of the intestinal lumen. Furthermore, absorption is accompanied by extensive conjugation and metabolism, and the forms appearing in the blood are usually different from the forms found in food. This indicates that *in vitro* experiments with the form of polyphenols found in food (the aglycone) are not necessarily relevant to the *in vivo* situation (8).

There are now intervention studies in the literature, of varying quality, that demonstrate significant biological effects of polyphenol consumption among humans, with the use of many different biomarkers (Tables 1–4). This review examines the effects demonstrated in some of the intervention studies reported in the literature. It considers most of the reports on quercetin, catechins, and procyanidins and some of those on isoflavones. Some of the reports described intervention studies involving consumption of foods and, in many of those cases, it was not proved that the observed effects were attributable to the polyphenol component. This situation may improve in the future, for example, with the use of isogenic lines of onions that differ only in their quercetin contents, allowing comparisons between groups consuming the same food but with different polyphenol contents. The bioavailability issues for each group of polyphenols are discussed in the context of the intervention studies.

HUMAN INTERVENTION STUDIES WITH FLAVONOIDS AND BIOAVAILABILITY ISSUES

Flavonols (including quercetin)

Quercetin is found at high concentrations in onions, apples, tea, broccoli, and red wine and as a component of *Ginkgo biloba*.

¹ From the Nutrient Bioavailability Group, Nestlé Research Center, Lausanne, Switzerland (GW), and the Unité des Maladies Métaboliques et Micronutriments, Institut National de la Recherche Agronomique, Saint-Genès Champanelle, France (CM).

² Presented at the 1st International Conference on Polyphenols and Health, held in Vichy, France, November 18–21, 2004.

³ Supported by Nestlé (Vevey, Switzerland).

⁴ Address reprint requests and correspondence to G Williamson, Nutrient Bioavailability Group, Nestlé Research Center, Vers chez les Blancs, 1000 Lausanne 26, Switzerland. E-mail: gary.williamson@rdls.nestle.com.

TABLE 1
Human intervention studies on flavonols or flavonol-containing foods¹

Substances given	Principal polyphenol ²	Dose per day	Days	No. of subjects per group ³	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
<i>Ginkgo biloba</i> extract EGb 761	Quercetin	120 mg extract	90	20	Decrease in blood pressure, increase in fasting plasma insulin and C-peptide	Liver function, coagulation tests, metabolic panel	9
<i>Ginkgo biloba</i> extract EGb 761	Quercetin	120 mg extract	90	6 hyperinsulinemic subjects	Enhanced hepatic extraction of insulin relative to C-peptide and reduction in plasma insulin	Plasma lipid profiles, blood cell counts	10
<i>Ginkgo biloba</i> extract EGb 761	Quercetin	160 mg extract	22	22 mountain climbers	Abolition of acute mountain sickness		11
<i>Ginkgo biloba</i>	Quercetin	320 mg extract	7	5 patients undergoing aortic valve replacement	Decrease in free radicals, delay in myoglobin release	Clinical improvement in outcome, but not significant	12
<i>Ginkgo biloba</i> extract EGb 761	Quercetin	320 mg extract	1	18 elderly subjects with slight age-related memory impairment	Increase in mental performance		13
"Phenol-rich diet"	Quercetin, kaempferol	21 mg quercetin, 9 mg kaempferol	6	19	Increase in erythrocyte superoxide dismutase activity, decrease in lymphocyte DNA damage (tail moment)	Plasma α -tocopherol and β -carotene	14
Onions	Quercetin	200 g onion	1	6	Increased resistance of lymphocyte DNA to strand breakage, decrease in urinary 8-hydroxy-2'-deoxyguanosine	Urinary malondialdehyde	15
Fried onions	Quercetin	50 mg	1	5	6% increase in plasma antioxidant capacity	Susceptibility of LDL to oxidation	16
Supplement	Quercetin	250 mg quercetin and other polyphenols	28	27		HDL/LDL cholesterol, platelet aggregation, plasma thromboxane B2, blood pressure, resting heart rate	17
Onions	Quercetin	114 mg	14	18		Platelet aggregation, thromboxane B2 production, other hemostatic variables	18
Parsley	Apigenin	114 mg	14	18		Platelet aggregation, thromboxane B2 production, other hemostatic variables	18
Onion and tea	Quercetin	76–110 mg quercetin and other flavonols	14	10 stable type 2 diabetic patients	Decrease in oxidative damage to lymphocyte DNA	Plasma vitamin C, α -tocopherol, urate, albumin, bilirubin, SOD, GPx, selenium	19
Supplement	Quercetin	30 mg	14	4	Decrease in TIMP-1 plasma protein and lymphocyte mRNA	TIMP-2 and matrix metalloprotein-2 lymphocyte mRNA or plasma protein	20
Supplement	Quercetin	100 mg [curcumin, 400 mg]	1	15 renal transplant patients	Improved renal function only in patients with elevated serum creatinine, improved urine output and lowered isoprostanes in patients with delayed graft function	Blood pressure, calcineurin levels	21

(Continued)

TABLE 1 (Continued)

Substances given	Principal polyphenol ²	Dose per day	Days	No. of subjects per group ³	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Supplement	Quercetin	2 × 500 mg	30	30 men with chronic pelvic pain syndrome	Improvement in NIH prostatitis symptom score		22
Supplement	Quercetin	30 mg	14	10	Improved oxidative resistance of LDL	Plasma triglycerides, total, HDL or LDL cholesterol, vitamin E or C, retinal and carotenoid	23
Supplement	Quercetin	2 × 500 mg	28	20	Improvement in cystitis symptoms	No side effects or adverse reactions	24

¹ SOD, superoxide dismutase; GPx, glutathione peroxidase; TIMP, tissue inhibitor of matrix metalloproteinase.

² The measured polyphenol; the dose is given in the next column. Ginkgo biloba also contains terpenes, and onions also contain high levels of biologically active sulfur compounds and other nutrients.

³ Healthy subjects, unless otherwise stated.

Intervention studies with quercetin are shown in Table 1. Some diverse effects have been demonstrated for *Ginkgo biloba*, but these may also be attributable to the terpenoid component of these extracts. Other studies have shown effects on antioxidant biomarkers, such as increased resistance of lymphocyte DNA to strand breakage, decreased urinary 8-hydroxy-2'-deoxyguanosine concentrations, increased plasma antioxidant capacity, decreased tissue inhibitor of matrix metalloproteinase-1 expression, altered renal function, improved prostatitis symptoms, and improved oxidative resistance to LDL. However, there are also studies that show no effects on these biomarkers.

Despite the lack of convincing evidence for consistent effects of quercetin in vivo in humans, there are numerous studies on the properties of quercetin in vitro. The apparent discrepancy between in vitro and in vivo studies may be partly attributable to absorption and metabolism. Generally, quercetin is not found in the plasma as the free form or as the parent glucoside. At the doses used in the intervention studies noted in Table 1 (21–1000 mg), it would be found exclusively as methyl, sulfate, or glucuronic acid conjugates (102); when added together, these compounds would represent the equivalent of ~1–5 $\mu\text{mol/L}$ aglycone equivalents at the highest dose. Lower doses of quercetin are more methylated than higher doses in humans (103). Quercetin has a relatively long plasma half-life of 11–28 h, and a 50-mg dose would lead to concentrations of up to ~0.75–1.5 $\mu\text{mol/L}$ in plasma (1, 4). There is only limited information on the properties of the conjugates in vitro, although it can be concluded that the conjugates have quantitatively different properties and are generally less biologically active, compared with the aglycones (104, 105). However, deconjugation could occur in some tissues such as the liver (106) or at sites of inflammation (107), leading to reactivation of the conjugated quercetin. The bioavailability issues may partially account for the lack of biological activity in vivo, although the lack of activity may reflect the short-term nature of the studies of quercetin and the selection of inappropriate biomarkers.

Isoflavones (genistein and daidzein)

The intervention studies on isoflavones are the most advanced and sophisticated of those for all of the polyphenols. Studies of the consumption of isoflavones lasting up to 1 year have shown effects on bone biomarkers, such as significant increases in bone mineral density and bone mineral content and changes in bone biomarkers, such as reduced excretion of pyridinium cross-links

and increased serum concentrations of bone-specific alkaline phosphatase and osteocalcin. Other effects include changes in LDL and HDL cholesterol concentrations, increases in LDL oxidation lag time, and changes in menopausal symptoms and hot flashes. Many, but not all, of the changes could be related to binding to the estrogen receptor, and this has been reviewed (108, 109).

Isoflavones occur in soy as glycosides, but some fermented products contain free aglycones (110). Consumption of isoflavone-rich foods or of the purified isoflavones themselves leads to appearance in the plasma, with a peak of absorption at 6–8 h (7). A dose of 50 mg of either daidzein or genistein, as typically used in intervention studies (the intervention studies noted in Table 1 used 37–128 mg per person per day), yields a peak plasma concentration of ~2 $\mu\text{mol/L}$ at ~6 h (4). The glycosides are not present in plasma, and most of the isoflavones are conjugated as sulfates or glucuronides; some free aglycone is also present, and Setchell et al (7) found 8% of the total daidzein as unconjugated aglycone 2 h after consumption of a dose of 50 mg. This decreased to 3% at steady state, which would apply to the intervention studies, because the studies were conducted for 14–365 d. After 4 wk of 30 mg/d isoflavones, the peak plasma concentrations showed no significant changes at the measurement times of 2 and 4 wk (111), which indicates that the bioavailability of isoflavones does not decrease during long periods of intake. However, we have unpublished data showing that isoflavone bioavailability increases during an extended period of intake (C Manach, unpublished observations, 2004), and this issue is currently unresolved. When administered in intervention studies, the isoflavones are clearly active and affect several biomarkers, especially related to bone. It is not known whether this effect is derived from free aglycone in the plasma, whether the conjugated forms are also active, or whether active deconjugation occurs in the relevant tissues. Some of the isoflavone conjugates are active in vitro (112), although information is very limited.

Catechins [(+)-catechin, (–)-epicatechin, (–)-epigallocatechin, (–)-epicatechin gallate, and (–)-epigallocatechin gallate]

(+)-Catechin and (–)-epicatechin are widely distributed in foods. Catechin concentrations are especially high in broad beans, black grapes, apricots, and strawberries. (–)-Epicatechin is found at high concentrations in apples, blackberries, broad

TABLE 2

Human intervention studies of isoflavones or isoflavone-containing foods

Substances given	Principal polyphenols ¹	Dose per day of isoflavones	Days	No. of subjects per group ²	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Soy foods	Genistein, daidzein	25–45 mg	30	6 female subjects (age: 20–29 y)	Follicular phase length increased, peak progesterone concentration delayed, midcycle peaks of luteinizing hormone and follicle-stimulating hormone suppressed, reduction in total cholesterol		25
Soy vs placebo	Genistein, daidzein	37 mg	70	20 postmenopausal women	Bone resorption	Bone stiffness	26
High, low, and control soy	Genistein, daidzein	73 mg	30	41 hypercholesterolemic men and postmenopausal women	Serum interleukin-6 increased in women	Serum C-reactive protein, serum amyloid A, serum α -tumor necrosis factor	27
Genistein	Genistein	54 mg	365	30 women (age: 47–57 y)	~3% increase in bone mineral density in femur and lumbar spine		28
Isoflavone concentrate	Genistein, daidzein	80 mg	14	29 healthy menopausal women		Endothelial function (flow-mediated dilation)	29
Isoflavone-rich soy	Genistein, daidzein	80 mg	168	24 perimenopausal women	Bone mineral density increased 5.6%, bone mineral content increased 10.1%		30
Isoflavone extract	Genistein, daidzein	100 mg	120	40 postmenopausal (2 y) women	Decrease in menopausal symptoms, decrease in total cholesterol and LDL cholesterol	Blood pressure, plasma glucose, HDL or triglyceride levels	31
Isoflavone-rich soy	Genistein, daidzein	56 mg	17	24	Plasma 8-epi-prostaglandin F2 decreased by ~20%, LDL oxidation lag time by 10%	LDL-tocopherol, plasma polyunsaturated fatty acids or malondialdehyde	32
Isoflavone tablet	Genistein, daidzein	80 mg	56	20 women (age: 50–70 y)		Blood pressure, plasma lipid, plasma lipoprotein, endothelium-(in)dependent dilation	33
Soy protein foods	Genistein, daidzein	86 mg	31	31 hyperlipidemic subjects	Decreased circulating oxidized LDL		34
Red clover	Genistein, daidzein, biochanin, formononetin	80 mg	35	17 women	Increase in arterial compliance by 23%	Plasma lipids	35
Soy protein	Isoflavones	90 mg	60	66 hypercholesterolemic postmenopausal women	Non-HDL cholesterol reduced, HDL cholesterol increased, mononuclear cell LDL receptor mRNA increased, increase in bone mineral content and density of lumbar spine		36
Soy protein powder	Isoflavones	128 mg	90	14 premenopausal women	Increase in plasma luteinizing hormone and follicle-stimulating hormone during preovulatory phase, decrease in free T ₃ and dehydroepiandrosterone sulfate during early follicular phase, decreased estrone during midfollicular phase	Length of menstrual cycle	37

(Continued)

TABLE 2 (Continued)

Substances given	Principal polyphenols ¹	Dose per day of isoflavones	Days	No. of subjects per group ²	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Genistein supplement	Genistein	54 mg	365	30 women (age: 47–57 y)	Reduced excretion of pyridinium cross-links, increased serum bone-specific alkaline phosphatase and osteocalcin, increase in bone mineral density in femur and lumbar spine		28
Supplement	Isoflavones	132 mg	84	32 postmenopausal women with diet-controlled type 2 diabetes mellitus	Lower fasting insulin, lower insulin resistance, lower LDL and total cholesterol, lower free thyroxine	HDL cholesterol, triglycerides, weight, blood pressure, creatinine, steroid hormones	38
Isoflavone-enriched soy protein	Isoflavones	90 mg	365	15 young healthy women		No changes in bone mineral density or other bone parameters	39
Supplement	Isoflavones	62 mg	28	23 healthy perimenopausal women	Reduced excretion of bone resorption biomarkers, LDL and serum cholesterol decreased	HDL or VLDL cholesterol, triglycerides, no liver cell damage and no tendency to diabetes	40

¹ The measured polyphenol; the dose is given in the next column.

² Healthy subjects, unless otherwise stated.

beans, cherries, black grapes, pears, raspberries, and chocolate. The gallates and the gallo catechins are found almost exclusively in tea, especially green tea. Deducing the intake of catechins during intervention studies is more difficult than for isoflavones, because the non-galloylated forms are widespread and can complicate intake estimates. This can lead to consumption of additional sources of catechins; furthermore, the amounts were not measured in some cases. The situation is clearer for the gallates or galloylated catechins, because they are almost exclusive to green tea. It is difficult to estimate the total flavonoid intake from black tea, because the fermentation process gives rise to a variety of structurally different oligomers. However, the amounts of remaining monomeric catechins can be readily estimated. Catechins are biologically active molecules that have a wide range of effects in vitro.

In human intervention studies (Table 3), catechins increased plasma antioxidant activity, as assessed with a variety of assays, decreased plasma lipid peroxide and malondialdehyde concentrations, increased plasma ascorbate concentrations, decreased nonheme iron absorption, and increased the resistance of LDL to oxidation. In addition, the green tea catechins, including the galloylated catechins, increased fat oxidation and energy expenditure and decreased the respiratory quotient and body weight. There were also some effects on vascular dilation.

The amounts of catechins administered in various intervention studies were highly variable, and administration was for short periods (1–28 d) (Table 1). With a dose of epigallocatechin gallate (EGCG) of 50 mg, peak plasma concentrations were ~0.15 $\mu\text{mol/L}$ (4). Although the exact percentages vary among individuals, among different studies, and with time after consumption, a substantial amount of the EGCG in plasma is unconjugated. For example, EGCG given to volunteers in one dose

of 2 mg/kg body wt yielded 77% of total EGCG as the unconjugated form at 1 h after consumption, with some individuals exhibiting values as high as 100% (113).

For the non-galloylated catechins, doses of 35 or 160 mg (+)-catechin (in red wine or chocolate) yielded plasma concentrations of 0.1–0.2 $\mu\text{mol/L}$ (6, 114), and a similar dose of (–)-epicatechin yielded 0.2 $\mu\text{mol/L}$ in plasma (114, 115). Both (+)-catechin and (–)-epicatechin are present in plasma exclusively as conjugates with methyl, sulfate, or glucuronic acid groups (6, 113, 116). Generally, catechins have short plasma half-lives (2–3 h).

In summary, the intervention studies with monomeric catechins give rise to plasma concentrations on the order of 0.1–0.5 $\mu\text{mol/L}$, but with rapid clearance. Substantial amounts of unconjugated forms of EGCG would be present in plasma, but all (+)-catechin and (–)-epicatechin is predicted to be conjugated. The percentage of catechins in the plasma that are sulfated or glucuronidated depends on the dose, but this is not usually measured in intervention studies.

Procyanidins [oligomeric (+)-catechin and (–)-epicatechin]

Procyanidins are oligomeric catechins, covalently linked together. Dimers and trimers are most common, but the degree of oligomerization can be quite high. Procyanidins are present at particularly high concentrations in cocoa, grapes/wine, and apples and are also found in many fruits, such as blackberries, cherries, figs, and plums (117). Purified procyanidins are weakly bioactive in vitro but exhibit numerous effects in vivo in intervention studies. It is important to note that procyanidins usually occur together with monomeric (+)-catechin and (–)-epicatechin; therefore, it is not clear whether the observed effects are

TABLE 3

Human intervention studies of monomeric catechins or (+)-catechin-containing foods¹

Substances given	Principal polyphenols ²	Dose per day	Days	No. of subjects per group ³	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Black tea, green tea, polyphenol-rich extract	Catechins	900 mL	28	13–16		Serum interleukin-6, interleukin-1 β , tumor necrosis factor- α , C-reactive protein, fibrinogen, plasminogen activator inhibitor-1, LDL oxidation, plasma cholesterol or triglycerides, plasma vitamin C or E	41, 42
Green tea drink	Catechins	5 g of green tea	1		Plasma ascorbic acid increased	Plasma β -carotene, α -tocopherol, uric acid	43
Green tea drink	Catechins	150, 300, 450 mL	1	10	Dose-dependent increase in plasma antioxidant activity		44
Green tea extract	Catechins	254 mg total catechins	1	18	40% decrease in plasma phospholipid hydroperoxide levels		45
Green tea, black tea	Catechins	8 cups	3	18		LDL oxidation ex vivo	46
Oolong tea	Catechins	0, 7.5, 15 g tea	3	12	Fat oxidation 12% higher		47
Green tea extract	Catechins	90 mg EGCG	1	10	Energy expenditure 4% higher, respiratory quotient 3% lower, 24-h urinary noradrenaline 40% higher		48
Green tea extract	Catechins	0.1 mmol	1	10	Decrease in nonheme iron absorption		49
Black tea	Catechins	5 cups	28	21	Endothelium-dependent dilation increased by 2.3%, independent dilation by 4.2%, lower soluble P-selectin	E-selectin, ICAM-1, VCAM-1, platelet aggregation, coagulation and fibrinolytic factors, F ₂ -isoprostane excretion	50–52
Green and black tea	Catechins	1 L	7	13	Trend to increasing ex vivo lag time for lipoprotein oxidation	F ₂ -isoprostane excretion	51, 53
Green tea extract	Catechins	375 mg catechins (270 mg EGCG)	90	70 Moderately obese patients	Body weight and waist circumference decreased		54
Green tea extract	Catechins	18 mg	21	8 smokers, 8 nonsmokers	Increased plasma antioxidant capacity	Plasma protein oxidation, plasma superoxide dismutase, glutathione peroxidase, catalase, urinary 8-oxodeoxyguanosine	55
Encapsulated green tea extract	Catechins	3 g extract (=10 cups)	28	20 female subjects on high-linoleic acid diet	Decrease in plasma malondialdehyde	Serum lipid, prostaglandins, nitric oxide metabolites, coagulation indicators	56
Green or black tea	Catechins	900 mL tea	28	45		Serum lipids, LDL oxidation	57
Black tea	Catechins	500 mL tea	1	10		Plasma antioxidant status	58
Black tea	Catechins	750 mL tea	28	14	Increase in LDL oxidation lag time ex vivo	Total cholesterol, triacylglycerol, apolipoprotein B	59

(Continued)

TABLE 3 (Continued)

Substances given	Principal polyphenols ²	Dose per day	Days	No. of subjects per group ³	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Black tea	Catechins	450 mL tea	1	49	Increased brachial dilation, not attributable to caffeine; small increase in systolic blood pressure	Nitroglycerin-induced endothelium-independent vasodilation, arterial diameter, diastolic blood pressure, heart rate, no effect on platelet aggregation	60, 61
Black tea	Catechins	900 mL	28	49	Increased brachial dilation, not attributable to caffeine	Nitroglycerin-induced, endothelium-independent vasodilation, arterial diameter, diastolic blood pressure or heart rate, no increase in systolic blood pressure; no effect on platelet aggregation	60, 61

¹ ICAM, intercellular adhesion molecule; VCAM, vascular cell adhesion molecule.

² The measured polyphenol; the dose is given in the next column.

³ Healthy subjects, unless otherwise stated.

attributable to the procyanidin component, the monomeric component, or both. The predominant effects are on the vascular system and include substantial increases in plasma antioxidant activity, decreased platelet aggregation (both stimulated and unstimulated), decreased plasma concentrations of lipid peroxide and thiobarbituric acid-reactive substances, decreased LDL cholesterol concentrations, increased HDL cholesterol concentrations, decreased susceptibility of LDL to oxidation, endothelium-dependent blood vessel dilation and decreased blood pressure, beneficial effects on capillary fragility and permeability, increased plasma ascorbate concentrations, decreased P-selectin expression, increased concentrations of nitrosated/nitrosylated species, decreased serum thromboxane concentrations, increased diameters of microvessels, reduced serum thromboxane B2 concentrations, increased plasma homocysteine concentrations, increased plasma vitamin B6 concentrations, maintenance of endothelial function (compared with loss with a high-fat diet), increased platelet-derived nitric oxide production, decreased superoxide release, increased α -tocopherol concentrations, and decreased concentrations of circulating autoantibodies to oxidized LDL (Table 4).

The metabolic fate of procyanidins after consumption is still a mystery. After consumption of 2 g of high-procyanidin grape seed extract by volunteers, the plasma concentrations of procyanidin B1 reached only 10 nmol/L (118); after consumption of 0.375 g cocoa/kg body wt, the plasma concentrations of procyanidin B2 reached only 41 nmol/L (114). When administered in a purified form to rats, procyanidin dimer B3 was not found in the plasma (119). However, human intervention studies with procyanidin-rich foods, as discussed above, show biological effects (Table 4). Either the effects are attributable to currently unidentified metabolites of the procyanidins or the effects are attributable to another component, such as the monomeric catechins (or both).

Microbial metabolites of polyphenols

The data on the bioavailability of polyphenols presented above considered only the presence of intact polyphenols in the blood,

ie, the ingested compound or its conjugates. The extensive microflora in the colon also plays a critical role in the metabolism of polyphenols. After microbial enzyme-catalyzed deconjugation of any polyphenol conjugates that reach the colon, there are 2 possible routes available, namely, absorption of the intact polyphenol through the colonic epithelium and passage into the bloodstream (as free or conjugated forms) or breakdown of the original polyphenol structure into metabolites. The absorption data presented above include the contribution of the absorption of intact polyphenols in the colon but do not include the breakdown contribution. Microbial metabolism deserves special consideration, because many of the diverse polyphenols are broken down into simpler phenolic compounds that are common to many different polyphenols. In addition, some of the microbial metabolites could have unique biological effects. For example, the isoflavone daidzein is converted to equol by gut microflora in ~30–40% of the population, and the equol is absorbed into the bloodstream in these people. There is emerging evidence that “equol producers” demonstrate better effects on some biomarkers, such as bone mineral density, after isoflavone consumption, compared with nonproducers (109). This is an example in which microflora activate a polyphenol to a more potent biologically active compound. Although intervention studies demonstrate an effect for procyanidins, the identity of the active component (or components) is not clear. Although intact procyanidins have some biological effects, they are poorly absorbed in an intact form (114, 118, 119). The active species could therefore be metabolites. Some low-molecular weight metabolites were identified in humans in vivo after consumption of cocoa procyanidins (120), but the biological activities of these metabolites are not known and remain to be investigated.

In summary, there are now many human intervention studies in the literature that show biological effects, but the exact effects depend on the class of polyphenol used. There are clear gaps. Most of the studies were short term, and there is a real need for longer-term studies; very few studies demonstrated a dose-response relationship, and this is also needed for convincing evidence. In addition, most studies, with the exception of those

TABLE 4

Human intervention studies of procyanidins or procyanidin-containing foods

Substances given	Principal polyphenols ¹	Dose per day	Days	No. of subjects per group ²	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Semisweet chocolate	Procyanidins, (-)-epicatechin	-420 mg procyanidins, 137 mg (-)-epicatechin	1	13	Increase in plasma antioxidant activity, decrease in plasma thiobarbituric acid-reactive substances		62
Chocolate chips	Procyanidins, catechins	220 mg flavonoids	1	18	Reduction in ADP/collagen-stimulated and adrenalin/collagen-stimulated, platelet-related primary hemostasis		63
Cocoa	Procyanidins, catechins	897 mg total	1	16	Inhibition of adrenalin-stimulated platelet activation and function		64
Cocoa powder/dark chocolate	Procyanidins, catechins	466 mg procyanidins	28	23	LDL oxidation lag time decreased by 8%, HDL cholesterol 4% higher	Thromboxane B2 or 6-keto-prostaglandin, plasma cholesterol, triglycerides	65
Cocoa	Procyanidins, catechins	70 mg (+)-catechin/(-)-epicatechin and 106 mg procyanidins (compared with equivalent with low polyphenols)	1	20	Endothelium-dependent dilation of brachial artery from 3.4% to 6.3%, plasma nitrosated/nitrosylated species increased by ~60%	Brachial artery diameter, forearm blood flow, blood pressure, heart rate, plasma nitrite, plasma nitrate	66
Cocoa tablets	Procyanidins, catechins	234 mg polyphenols	28	32	Increase in plasma ascorbic acid, lower P-selectin expression, lower ADP-induced platelet aggregation, lower collagen-induced platelet aggregation	Plasma antioxidant status, plasma oxidation biomarkers	67
Cocoa	Procyanidins, catechins	500 mg polyphenols	14	13	Lower systolic and diastolic blood pressure	Heart rate, plasma cholesterol, triglyceride, and glucose concentrations	68
Chocolate	Procyanidins, catechins	320 mg procyanidin, 104 mg (-)-epicatechin	1	20	Small increase in plasma antioxidant activity	Plasma 8-isoprostane	69
Cocoa	Procyanidins, catechins	100 g dark chocolate	1	12	Increase in plasma antioxidant activity by 20%	Addition of milk reduced the effect on plasma antioxidant activity	70
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	75 mg	30	30 women with melasma	Decrease in average melasma area		71
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	150 mg	42	25	Increase of plasma oxygen radical absorbance capacity, decrease in LDL cholesterol, increase in HDL cholesterol	LDL oxidizability, plasma lipid peroxides	72
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	200 mg	56	11	Decrease in systolic blood pressure, decrease in serum thromboxane		73
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	120 mg for 30 d, then 60 mg for 30 d	60	11	Decrease in serum reactive oxygen species, lymphocyte apoptosis and p56 ^{lck} , reduction in erythrocyte sedimentation rate		74

(Continued)

TABLE 4 (Continued)

Substances given	Principal polyphenols ¹	Dose per day	Days	No. of subjects per group ²	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	300 mg	60	40 with chronic venous insufficiency	Disappearance of edema and pain, reduction in leg heaviness	Venous blood flow	75
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	100 mg	1 dose	16 smokers	Decrease in smoking-induced platelet reactivity		76
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	150 mg	28	60	Decreased platelet aggregation, increased diameter of microvessels	Blood glucose and lipids	77
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	200 mg	56	19 smokers	Reduced platelet activity, reduced serum thromboxane B2	(No effect in non-smokers)	78
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	200 mg	30	24 athletes	Increased endurance		79
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	200 mg	90	4 sub-fertile men	Improved sperm morphologic features	Ham's F10 capacitated count, motility score	80
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	1.1, then 1.7 mg/kg body weight	28 + 28	21	Decrease in ultraviolet light sensitivity		81
Pycnogenol (French pine bark extract)	Procyanidins, some (+)-catechin	30 mg	14 + 28	39 patients with endometriosis, severe menstrual pain or chronic pelvic pain	Reduction in menstrual cramps, abdominal pain, and tenderness		82
Cranberry juice	Procyanidins	50 mL juice	180	150 women with urinary tract infections	20% decrease in recurrence of urinary tract infections		83
Cranberry juice	Procyanidins	500 mL juice	1 dose	9	Plasma vitamin C increased by 30%, plasma antioxidant activity increased		84
Cranberry juice	Procyanidins	354 mL juice		112 men with prostate cancer undergoing radiotherapy		Urinary symptoms	85
Blueberry	Procyanidins, anthocyanins, quercetin	500 mL juice	1	9		Plasma vitamin C, plasma antioxidant activity	84
Black currant and apple juice	Procyanidins, anthocyanins, quercetin	1.5 L (9.6 mg quercetin)	5	7	Plasma malondialdehyde decreased, plasma 2-amino-adipic acid semialdehyde increased	Erythrocyte 2-aminoadipic semialdehyde, plasma trolox equivalent antioxidant activity, γ -glutamyl semialdehyde	86
Red wine	Procyanidins, anthocyanins, quercetin	500 mL	1	5		No effect on immune functions (tumor necrosis factor- α , interleukin-2, interleukin-4, lymphocyte proliferation, phagocytic activity)	87
Red wine	Procyanidins, anthocyanins, quercetin	375 mL	14	9	Decreased plasma lipid peroxides	LDL or HDL cholesterol, plasma triacylglycerol	88
Red wine	Procyanidins, anthocyanins, quercetin	4 glasses	21	11	Increased plasma homocysteine, increased plasma vitamin B6		89
Red wine	Procyanidins, anthocyanins, quercetin	240 mL wine	30	6	Maintenance of endothelial function, compared with loss with a high-fat diet		90

(Continued)

TABLE 4 (Continued)

Substances given	Principal polyphenols ¹	Dose per day	Days	No. of subjects per group ²	Biomarkers significantly affected	Biomarkers not significantly affected	Ref
Purple grape juice	Procyanidins	7 mL/kg	14	20	Platelet aggregation inhibited, platelet-derived nitric oxide production increased, superoxide release decreased, α -tocopherol increased, plasma protein-independent activity increased		91
Purple grape juice	Procyanidins	2 × 8mL/kg body weight	56	22 (age: 64 ± 10 y) with severe endothelial dysfunction	Increased flow-mediated dilation of the brachial artery	Plasma glucose or insulin, nitroglycerin-mediated vasodilation	92
Grape juice	Procyanidins	400 mL juice	7–10	10	Platelet aggregation reduced by 77%	No effect of orange or grapefruit juice	93
Grape polyphenol extract	Procyanidins	300 mg	28	24 male smokers	Small decrease in plasma thiobarbituric acid-reactive substances and increase in LDL oxidation lag phase	HDL, LDL, or total cholesterol	94
Grape seed extract	Procyanidins	100 mg	60	40 hypercholesterolemic subjects	Decrease in circulating auto-antibodies to oxidized LDL	LDL/HDL cholesterol, triglycerides	95
Purple grape juice	Procyanidins	7.7 mL/kg	14	15 adults with coronary artery disease	Flow-mediated dilation of blood vessels, LDL lag time increased		96
Pomegranate juice	Polyphenol mixture	1.5 mmol total polyphenols	14	10 hypertensive patients	Increased serum paraoxonase activity, increased plasma total antioxidant activity, increase in resistance of LDL to oxidation, decreased serum angiotensin-converting enzyme, decreased cytochrome P450 activity, decreased LDL susceptibility to aggregation, increased serum paraoxonase	Total cholesterol, LDL cholesterol, triglycerides	97, 98
Endothelon ³	Procyanidins	100 mg	14	37 elderly subjects with microcirculatory problems	Beneficial effect on capillary fragility and permeability		99
Endothelon ³	Procyanidins	200 mg	35	50	Improvement of the visual performance after glare, of the visual adaptation to low luminances		100
Endothelon ³	Procyanidins	150 mg	30–60	13 hypertensive and diabetic patients	Increase of capillary resistance		101

¹ The measured polyphenol; the dose is given in the next column.

² Healthy subjects, unless otherwise stated.

³ Endothelon, standardized grape seed extract.

with isoflavones, administered food instead of pure compounds, and the effects noted may thus be attributable to some other component in that food. Finally, metabolism by microflora needs to be understood, because the gut microflora probably plays a major role in the biological activity of many polyphenols. ☞

REFERENCES

- Scalbert A, Williamson G. Dietary intake and bioavailability of polyphenols. *J Nutr* 2000;130:2073S–85S.
- Rowland I, Faughnan M, Hoey L, Wahala K, Williamson G, Cassidy A. Bioavailability of phyto-oestrogens. *Br J Nutr* 2003;89(suppl 1): 838–52.
- Day AJ, Williamson G. Biomarkers for exposure to dietary flavonoids: a review of the current evidence for identification of quercetin glycosides in plasma. *Br J Nutr* 2001;86:S105–10.
- Manach C, Williamson G, Morand C, Scalbert A, Rémésy C. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* 2005;81(suppl):230S–42S.
- Hollman PC, van Trijp JM, Buysman MN, et al. Relative bioavailability

- of the antioxidant flavonoid quercetin from various foods in man. *FEBS Lett* 1997;418:152–6.
6. Donovan JL, Bell JR, Kasim-Karakas S, et al. Catechin is present as metabolites in human plasma after consumption of red wine. *J Nutr* 1999;129:1662–8.
 7. Setchell KD, Brown NM, Desai P, et al. Bioavailability of pure isoflavones in healthy humans and analysis of commercial soy isoflavone supplements. *J Nutr* 2001;131:1362S–75S.
 8. Williamson G. The use of flavonoid aglycones in in vitro systems to test biological activities: based on bioavailability data, is this a valid approach? *Phytochem Rev* 2003;1:215–222.
 9. Kudolo GB. The effect of 3-month ingestion of *Ginkgo biloba* extract on pancreatic β -cell function in response to glucose loading in normal glucose-tolerant individuals. *J Clin Pharmacol* 2000;40:647–54.
 10. Kudolo GB. The effect of 3-month ingestion of *Ginkgo biloba* extract (EGb 761) on pancreatic β -cell function in response to glucose loading in individuals with non-insulin-dependent diabetes mellitus. *J Clin Pharmacol* 2001;41:600–11.
 11. Roncin JP, Schwartz F, d'Arbigny P. EGb 761 in control of acute mountain sickness and vascular reactivity to cold exposure. *Aviat Space Environ Med* 1996;67:445–52.
 12. Pietri S, Seguin JR, d'Arbigny P, Drieu K, Culcasi M. *Ginkgo biloba* extract (EGb 761) pretreatment limits free radical-induced oxidative stress in patients undergoing coronary bypass surgery. *Cardiovasc Drugs Ther* 1997;11:121–31.
 13. Allain H, Raoul P, Lieury A, LeCoz F, Gandon JM, d'Arbigny P. Effect of two doses of *Ginkgo biloba* extract (EGb 761) on the dual-coding test in elderly subjects. *Clin Ther* 1993;15:549–58.
 14. Kim HY, Kim OH, Sung MK. Effects of phenol-depleted and phenol-rich diets on blood markers of oxidative stress, and urinary excretion of quercetin and kaempferol in healthy volunteers. *J Am Coll Nutr* 2003;22:217–23.
 15. Boyle SP, Dobson VL, Duthie SJ, Kyle JAM, Collins AR. Absorption and DNA protective effects of flavonoid glycosides from an onion meal. *Eur J Nutr* 2000;39:213–23.
 16. McAnlis GT, Mceneny J, Pearce J, Young IS. Absorption and antioxidant effects of quercetin from onions, in man. *Eur J Clin Nutr* 1999;53:92–6.
 17. Conquer JA, Mainai G, Azzini E, Raguzzini A, Holub BJ. Supplementation with quercetin markedly increases plasma quercetin concentration without effect on selected risk factors for heart disease in healthy subjects. *J Nutr* 1998;128:593–7.
 18. Janssen PLTM, Mensink RP, Cox FJJ, et al. Effects of the flavonoids quercetin and apigenin on hemostasis in healthy volunteers: results from an in vitro and a dietary supplement study. *Am J Clin Nutr* 1998;67:255–62.
 19. Lean MEJ, Noroozi M, Kelly I, et al. Dietary flavonols protect diabetic human lymphocytes against oxidative damage to DNA. *Diabetes* 1999;48:176–81.
 20. Morrow DM, Fitzsimmons PE, Chopra M, McGlynn H. Dietary supplementation with the anti-tumour promoter quercetin: its effects on matrix metalloproteinase gene regulation. *Mutat Res* 2001;480–481:269–76.
 21. Shoskes DA, Thomas M, Pobgee R, Fromkin B, Copley JB. Phase I study of oral bioflavonoids in cadaveric renal transplant recipients: effects on delayed graft function and calcineurin inhibitor toxicities. *Transplant Proc* 2003;35:841–2.
 22. Shoskes DA, Zeitlin SI, Shahed A, Rajfer J. Quercetin in men with category III chronic prostatitis: a preliminary prospective, double-blind, placebo-controlled trial. *Urology* 1999;54:960–3.
 23. Chopra M, Fitzsimmons PE, Strain JJ, Thurnham DI, Howard AN. Non-alcoholic red wine extract and quercetin inhibit LDL oxidation without affecting plasma antioxidant vitamin and carotenoid concentrations. *Clin Chem* 2000;46:1162–70.
 24. Katske F, Shoskes DA, Sender M, Poliakin R, Gagliano K, Rajfer J. Treatment of interstitial cystitis with a quercetin supplement. *Tech Urol* 2001;7:44–6.
 25. Cassidy A, Bingham S, Setchell K. Biological effects of isoflavones in young women: importance of the chemical composition of soyabean products. *Br J Nutr* 1995;74:587–601.
 26. Yamori Y, Moriguchi EH, Teramoto T, et al. Soybean isoflavones reduce postmenopausal bone resorption in female Japanese immigrants in Brazil: a ten-week study. *J Am Coll Nutr* 2002;21:560–3.
 27. Jenkins DJ, Kendall CW, Connelly PW, et al. Effects of high- and low-isoflavone (phytoestrogen) soy foods on inflammatory biomarkers and proinflammatory cytokines in middle-aged men and women. *Metabolism* 2002;51:919–24.
 28. Morabito N, Crisafulli A, Vergara C, et al. Effects of genistein and hormone-replacement therapy on bone loss in early postmenopausal women: a randomized double-blind placebo-controlled study. *J Bone Miner Res* 2002;17:1904–12.
 29. Hale G, Paul-Labrador M, Dwyer JH, Merz CN. Isoflavone supplementation and endothelial function in menopausal women. *Clin Endocrinol* 2002;56:693–701.
 30. Alekel DL, Germain AS, Peterson CT, Hanson KB, Stewart JW, Toda T. Isoflavone-rich soy protein isolate attenuates bone loss in the lumbar spine of perimenopausal women. *Am J Clin Nutr* 2000;72:844–52.
 31. Han KK, Soares JM Jr, Haidar MA, de Lima GR, Baracat EC. Benefits of soy isoflavone therapeutic regimen on menopausal symptoms. *Obstet Gynecol* 2002;99:389–94.
 32. Wiseman H, O'Reilly JD, Adlercreutz H, et al. Isoflavone phytoestrogens consumed in soy decrease F_2 -isoprostane concentrations and increase resistance of low-density lipoprotein to oxidation in humans. *Am J Clin Nutr* 2000;72:395–400.
 33. Simons LA, von Konigsmark M, Simons J, Celermajer DS. Phytoestrogens do not influence lipoprotein levels or endothelial function in healthy, postmenopausal women. *Am J Cardiol* 2000;85:1297–301.
 34. Jenkins DJ, Kendall CW, Garsetti M, et al. Effect of soy protein foods on low-density lipoprotein oxidation and ex vivo sex hormone receptor activity: a controlled crossover trial. *Metabolism* 2000;49:537–43.
 35. Nestel PJ, Pomeroy S, Kay S, et al. Isoflavones from red clover improve systemic arterial compliance but not plasma lipids in menopausal women. *J Clin Endocrinol Metab* 1999;84:895–8.
 36. Potter SM, Baum JA, Teng H, Stillman RJ, Shay NF, Erdman JW Jr. Soy protein and isoflavones: their effects on blood lipids and bone density in postmenopausal women. *Am J Clin Nutr* 1998;68:1375S–9S.
 37. Duncan AM, Merz BE, Xu X, Nagel TC, Phipps WR, Kurzer MS. Soy isoflavones exert modest hormonal effects in premenopausal women. *J Clin Endocrinol Metab* 1999;84:192–7.
 38. Jayagopal V, Albertazzi P, Kilpatrick ES, et al. Beneficial effects of soy phytoestrogen intake in postmenopausal women with type 2 diabetes. *Diabetes Care* 2002;25:1709–14.
 39. Anderson JJ, Chen X, Boass A, et al. Soy isoflavones: no effects on bone mineral content and bone mineral density in healthy, menstruating young adult women after one year. *J Am Coll Nutr* 2002;21:388–93.
 40. Uesugi T, Fukui Y, Yamori Y. Beneficial effects of soybean isoflavone supplementation on bone metabolism and serum lipids in postmenopausal Japanese women: a four-week study. *J Am Coll Nutr* 2002;21:97–102.
 41. de Maat MP, Pijl H, Klufft C, Princen HM. Consumption of black and green tea had no effect on inflammation, haemostasis and endothelial markers in smoking healthy individuals. *Eur J Clin Nutr* 2000;54:757–63.
 42. Princen HM, van Duyvenvoorde W, Buytenhek R, et al. No effect of consumption of green and black tea on plasma lipid and antioxidant levels and on LDL oxidation in smokers. *Arterioscler Thromb Vasc Biol* 1998;18:833–41.
 43. Gomikawa S, Ishikawa Y. Effects of catechins and ground green tea drinking on the susceptibility of plasma and LDL to the oxidation in vitro and ex vivo. *J Clin Biochem Nutr* 2002;32:55–68.
 44. Sung H, Nah J, Chun S, Park H, Yang SE, Min WK. In vivo antioxidant effect of green tea. *Eur J Clin Nutr* 2000;54:527–9.
 45. Nakagawa K, Ninomiya M, Okubo T, et al. Tea catechin supplementation increases antioxidant capacity and prevents phospholipid hydroperoxidation in plasma of humans. *J Agric Food Chem* 1999;47:3967–73.
 46. het Hof KH, Wiseman SA, Yang CS, Tijburg LB. Plasma and lipoprotein levels of tea catechins following repeated tea consumption. *Proc Soc Exp Biol Med* 1999;220:203–9.
 47. Rumpler W, Seale J, Clevidence B, et al. Oolong tea increases metabolic rate and fat oxidation in men. *J Nutr* 2001;131:2848–52.
 48. Dulloo AG, Duret C, Rohrer D, et al. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *Am J Clin Nutr* 1999;70:1040–5.
 49. Samman S, Sandstrom B, Toft MB, et al. Green tea or rosemary extract added to foods reduces nonheme-iron absorption. *Am J Clin Nutr* 2001;73:607–12.
 50. Hodgson JM, Puddey IB, Burke V, Watts GF, Beilin LJ. Regular

- ingestion of black tea improves brachial artery vasodilator function. *Clin Sci* 2002;102:195–201.
51. Hodgson JM, Croft KD, Mori TA, Burke V, Beilin LJ, Puddey IB. Regular ingestion of tea does not inhibit in vivo lipid peroxidation in humans. *J Nutr* 2002;132:55–8.
 52. Hodgson JM, Puddey IB, Mori TA, Burke V, Baker RI, Beilin LJ. Effects of regular ingestion of black tea on haemostasis and cell adhesion molecules in humans. *Eur J Clin Nutr* 2001;55:881–6.
 53. Hodgson JM, Puddey IB, Croft KD, et al. Acute effects of ingestion of black and green tea on lipoprotein oxidation. *Am J Clin Nutr* 2000;71:1103–7.
 54. Chantre P, Lairon D. Recent findings of green tea extract AR25 (Exolise) and its activity for the treatment of obesity. *Phytotherapy* 2002;9:3–8.
 55. Young JF, Dragstedt LO, Haraldsdottir J, et al. Green tea extract only affects markers of oxidative status postprandially: lasting antioxidant effect of flavonoid-free diet. *Br J Nutr* 2002;87:343–55.
 56. Freese R, Basu S, Hietanen E, et al. Green tea extract decreases plasma malondialdehyde concentration but does not affect other indicators of oxidative stress, nitric oxide production, or hemostatic factors during a high-linoleic acid diet in healthy females. *Eur J Nutr* 1999;38:149–57.
 57. het Hof KH, de Boer HS, Wiseman SA, Lien N, Westrate JA, Tijburg LB. Consumption of green or black tea does not increase resistance of low-density lipoprotein to oxidation in humans. *Am J Clin Nutr* 1997;66:1125–32.
 58. Maxwell S, Thorpe G. Tea flavonoids have little short term impact on serum antioxidant activity. *Br Med J* 1996;313:229.
 59. Ishikawa T, Suzukawa M, Ito T, et al. Effect of tea flavonoid supplementation on the susceptibility of low-density lipoprotein to oxidative modification. *Am J Clin Nutr* 1997;66:261–6.
 60. Duffy SJ, Keaney JF Jr, Holbrook M, et al. Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation* 2001;104:151–6.
 61. Duffy SJ, Vita JA, Holbrook M, Swerdloff PL, Keaney JF Jr. Effect of acute and chronic tea consumption on platelet aggregation in patients with coronary artery disease. *Arterioscler Thromb Vasc Biol* 2001;21:1084–9.
 62. Rein D, Lotito S, Holt RR, Keen CL, Schmitz HH, Fraga CG. Epicatechin in human plasma: in vivo determination and effect of chocolate consumption on plasma oxidation status. *J Nutr* 2000;130:2109S–14S.
 63. Holt RR, Schramm DD, Keen CL, Lazarus SA, Schmitz HH. Chocolate consumption and platelet function. *JAMA* 2002;287:2212–3.
 64. Pearson DA, Paglieroni TG, Rein D, et al. The effects of flavanol-rich cocoa and aspirin on ex vivo platelet function. *Thromb Res* 2002;106:191–7.
 65. Wan Y, Vinson JA, Etherton TD, Proch J, Lazarus SA, Kris-Etherton PM. Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and prostaglandin concentrations in humans. *Am J Clin Nutr* 2001;74:596–602.
 66. Heiss C, Dejam A, Kleinbongard P, Schewe T, Sies H, Kelm M. Vascular effects of cocoa rich in flavan-3-ols. *JAMA* 2003;290:1030–1.
 67. Murphy KJ, Chronopoulos AK, Singh I, et al. Dietary flavanols and procyanidin oligomers from cocoa (*Theobroma cacao*) inhibit platelet function. *Am J Clin Nutr* 2003;77:1466–73.
 68. Taubert D, Berkels R, Roosen R, Klaus W. Chocolate and blood pressure in elderly individuals with isolated systolic hypertension. *JAMA* 2003;290:1029–30.
 69. Wang JF, Schramm DD, Holt RR, et al. A dose-response effect from chocolate consumption on plasma epicatechin and oxidative damage. *J Nutr* 2000;130:2115S–9S.
 70. Serafini M, Bugianesi R, Maiani G, Valtuena S, De Santis S, Crozier A. Plasma antioxidants from chocolate. *Nature* 2003;424:1013.
 71. Ni Z, Mu Y, Gulati O. Treatment of melasma with Pycnogenol. *Phytother Res* 2002;16:567–71.
 72. Devaraj S, Vega-Lopez S, Kaul N, Schonlau F, Rohdewald P, Jialal I. Supplementation with a pine bark extract rich in polyphenols increases plasma antioxidant capacity and alters the plasma lipoprotein profile. *Lipids* 2002;37:931–4.
 73. Hosseini S, Lee J, Sepulveda RT, Rohdewald P, Watson RR. A randomized, double-blind, placebo-controlled, prospective, 16-week crossover study to determine the role of Pycnogenol in modifying blood pressure in mildly hypertensive patients. *Nutr Res* 2001;21:1251–60.
 74. Stefanescu M, Matache C, Onu A, et al. Pycnogenol efficacy in the treatment of systemic lupus erythematosus patients. *Phytother Res* 2001;15:698–704.
 75. Arcangeli P. Pycnogenol in chronic venous insufficiency. *Fitoterapia* 2000;71:236–44.
 76. Putter M, Grotemeyer KHM, Wurthwein G, et al. Inhibition of smoking-induced platelet aggregation by aspirin and Pycnogenol. *Thromb Res* 1999;95:155–61.
 77. Wang S, Tan D, Zhao Y, Gao G, Gao X, Hu L. The effect of Pycnogenol on the microcirculation, platelet function and ischemic myocardium in patients with coronary artery diseases. *Eur Bull Drug Res* 1999;7:19–25.
 78. Araghi-Niknam M, Hosseini S, Larson D, Rohdewald P, Watson RR. Pine bark extract reduces platelet aggregation. *Integr Med* 1999;2:73–7.
 79. Pavlovic P. Improved endurance by use of antioxidants. *Eur Bull Drug Res* 1999;7:26–9.
 80. Roseff SJ. Improvement in sperm quality and function with French maritime pine tree bark extract. *J Reprod Med* 2002;47:821–4.
 81. Saliou C, Rimbach G, Moini H, et al. Solar ultraviolet-induced erythema in human skin and nuclear factor- κ B-dependent gene expression in keratinocytes are modulated by a French maritime pine bark extract. *Free Radic Biol Med* 2001;30:154–60.
 82. Kohama T, Suzuki N. The treatment of gynaecological disorders with Pycnogenol. *Eur Bull Drug Res* 1999;7:30–2.
 83. Kontiokari T, Sundqvist K, Nuutinen M, Pokka T, Koskela M, Uhari M. Randomised trial of cranberry-lingonberry juice and *Lactobacillus* GG drink for the prevention of urinary tract infections in women. *Br Med J* 2001;322:1571.
 84. Pedersen CB, Kyle J, Jenkinson AM, Gardner PT, McPhail DB, Duthie GG. Effects of blueberry and cranberry juice consumption on the plasma antioxidant capacity of healthy female volunteers. *Eur J Clin Nutr* 2000;54:405–8.
 85. Campbell G, Pickles T, D'yachkova Y. A randomised trial of cranberry versus apple juice in the management of urinary symptoms during external beam radiation therapy for prostate cancer. *Clin Oncol* 2003;15:322–8.
 86. Young JF, Nielsen SE, Haraldsdottir J, et al. Effect of fruit juice intake on urinary quercetin excretion and biomarkers of antioxidative status. *Am J Clin Nutr* 1999;69:87–94.
 87. Watzl B, Bub A, Briviba K, Rechkemmer G. Acute intake of moderate amounts of red wine or alcohol has no effect on the immune system of healthy men. *Eur J Nutr* 2002;41:264–70.
 88. Nigdikar SV, Williams NR, Griffin BA, Howard AN. Consumption of red wine polyphenols reduces the susceptibility of low-density lipoproteins to oxidation in vivo. *Am J Clin Nutr* 1998;68:258–65.
 89. van der Gaag MS, Ubbink JB, Sillanaukee P, Nikkari S, Hendriks HF. Effect of consumption of red wine, spirits, and beer on serum homocysteine. *Lancet* 2000;355:1522.
 90. Cuevas AM, Guasch V, Castillo O, et al. A high-fat diet induces and red wine counteracts endothelial dysfunction in human volunteers. *Lipids* 2000;35:143–8.
 91. Freedman JE, Parker C, III, Li L, et al. Select flavonoids and whole juice from purple grapes inhibit platelet function and enhance nitric oxide release. *Circulation* 2001;103:2792–8.
 92. Chou EJ, Keevil JG, Aeschlimann S, Wiebe DA, Folts JD, Stein JH. Effect of ingestion of purple grape juice on endothelial function in patients with coronary heart disease. *Am J Cardiol* 2001;88:553–5.
 93. Keevil JG, Osman HE, Reed JD, Folts JD. Grape juice, but not orange juice or grapefruit juice, inhibits human platelet aggregation. *J Nutr* 2000;130:53–6.
 94. Vigna GB, Costantini F, Aldini G, et al. Effect of a standardized grape seed extract on low-density lipoprotein susceptibility to oxidation in heavy smokers. *Metabolism* 2003;52:1250–7.
 95. Preuss HG, Wallerstedt D, Talpur N, et al. Effects of niacin-bound chromium and grape seed proanthocyanidin extract on the lipid profile of hypercholesterolemic subjects: a pilot study. *J Med* 2000;31:227–46.
 96. Stein JH, Keevil JG, Wiebe DA, Aeschlimann S, Folts JD. Purple grape juice improves endothelial function and reduces the susceptibility of LDL cholesterol to oxidation in patients with coronary artery disease. *Circulation* 1999;100:1050–5.
 97. Aviram M, Dornfeld L, Kaplan M, et al. Pomegranate juice flavonoids inhibit low-density lipoprotein oxidation and cardiovascular diseases:



- studies in atherosclerotic mice and in humans. *Drugs Exp Clin Res* 2002;28:49–62.
98. Aviram M, Dornfeld L, Rosenblat M, et al. Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL, and platelet aggregation: studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *Am J Clin Nutr* 2000;71:1062–76.
 99. Dartenuc JY, Marache P, Choussat H. Résistance capillaire en gériatrie etude d'un microangioprotecteur: endotélon. (Hair resistance in geriatrics studied with a microangioprotector: endothelion.) *Bordeaux Med* 1980;13:903–7 (in French).
 100. Corbé C, Boissin JP, Siou A. Sens lumineux et circulation chorioretinienne: etude de l'effet des O.P.C. (Endotélon). (Luminosity and chorioretic circulation: a study on the effects of O.P.C. (endothelion).) *J Fr Ophthalmol* 1998;11:453–60 (in French).
 101. Lagrue G, Olivier-Martin F, Grillot A. A study of the effects of procyanidin oligomers on capillary resistance in hypertension and in certain nephropathies. *Sem Hop* 1981;57:1399–401 (in French).
 102. Day AJ, Mellon FA, Barron D, Sarrazin G, Morgan MRA, Williamson G. Human metabolism of dietary flavonoids: identification of plasma metabolites of quercetin. *Free Radic Res* 2001;212:941–52.
 103. DuPont MS, Mondin Z, Williamson G, Price KR. Effect of variety, processing, and storage on the flavonoid glycoside content and composition of lettuce and endive. *J Agric Food Chem* 2000;48:3957–64.
 104. Day AJ, Bao Y, Morgan MRA, Williamson G. Conjugation position of quercetin glucuronides and effect on biological activity. *Free Radic Biol Med* 2000;29:1234–43.
 105. Cano A, Arnao MB, Williamson G, Garcia-Conesa MT. Superoxide scavenging by polyphenols: effect of conjugation and dimerization. *Redox Rep* 2002;7:379–83.
 106. O'Leary KA, Day AJ, Needs P, Mellon FA, O'Brien NM, Williamson G. Metabolism of quercetin-7- and quercetin-3-glucuronides by an in vitro hepatic model: the role of human β -glucuronidase, sulfotransferase, catechol-O-methyltransferase and multi-resistant protein 2 (MRP2) in flavonoid metabolism. *Biochem Pharmacol* 2003;65:479–91.
 107. Shimoi K, Saka N, Nozawa R, et al. Deglucuronidation of a flavonoid, luteolin monoglucuronide, during inflammation. *Drug Metab Dispos* 2001;29:1521–4.
 108. Setchell KD, Cassidy A. Dietary isoflavones: biological effects and relevance to human health. *J Nutr* 1999;129:758S–67S.
 109. Setchell KD, Brown NM, Lydeking-Olsen E. The clinical importance of the metabolite equol: a clue to the effectiveness of soy and its isoflavones. *J Nutr* 2002;132:3577–84.
 110. Kishida T, Ataki H, Takebe M, Ebihara K. Soybean meal fermented by *Aspergillus awamori* increases the cytochrome P-450 content of the liver microsomes of mice. *J Agric Food Chem* 2000;48:1367–72.
 111. Izumi T, Piskula MK, Osawa S, et al. Soy isoflavone aglycones are absorbed faster and in higher amounts than their glucosides in humans. *J Nutr* 2000;130:1695–9.
 112. Zhang Y, Song TT, Cunnick JE, Murphy PA, Hendrich S. Daidzein and genistein glucuronides in vitro are weakly estrogenic and activate human natural killer cells at nutritionally relevant concentrations. *J Nutr* 1999;129:399–405.
 113. Lee MJ, Maliakal P, Chen LS, et al. Pharmacokinetics of tea catechins after ingestion of green tea and (–)-epigallocatechin-3-gallate by humans: formation of different metabolites and individual variability. *Cancer Epidemiol Biomarkers Prev* 2002;11:1025–32.
 114. Holt RR, Lazarus SA, Sullards MC, et al. Procyanidin dimer B2 [epicatechin-(4 β -8)-epicatechin] in human plasma after the consumption of a flavanol-rich cocoa. *Am J Clin Nutr* 2002;76:798–804.
 115. Yang CS, Chen LS, Lee MJ, Balentine D, Kuo MC, Schantz SP. Blood and urine levels of tea catechins after ingestion of different amounts of green tea by human volunteers. *Cancer Epidemiol Biomarkers Prev* 1998;7:351–4.
 116. Bell JR, Donovan JL, Wong R, et al. (+)-Catechin in human plasma after ingestion of a single serving of reconstituted red wine. *Am J Clin Nutr* 2000;71:103–8.
 117. Pascual-Teresa S, Santos-Buelga C, Rivas-Gonzalo JC. Quantitative analysis of flavan-3-ols in Spanish foodstuffs and beverages. *J Agric Food Chem* 2000;48:5331–7.
 118. Sano A, Yamakoshi J, Tokutake S, Tobe K, Kubota Y, Kikuchi M. Procyanidin B1 is detected in human serum after intake of proanthocyanidin-rich grape seed extract. *Biosci Biotechnol Biochem* 2003;67:1140–3.
 119. Donovan JL, Manach C, Rios L, Morand C, Scalbert A, Remesy C. Procyanidins are not bioavailable in rats fed a single meal containing a grape seed extract or the procyanidin dimer B-3. *Br J Nutr* 2002;87:299–306.
 120. Rios LY, Gonthier MP, Remesy C, et al. Chocolate intake increases urinary excretion of polyphenol-derived phenolic acids in healthy human subjects. *Am J Clin Nutr* 2003;77:912–8.

